Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

County: 001

Page:

Run Date:

6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|----------------------|----------------------------------|--------|----------------------|
| | | Anemia Low Hb/Hct | | | Anemia Low Hb/Hct |
| | Number | % | | Number | % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 21 | * | 6 - 11 Months | 39 | * |
| 12 - 17 Months | 49 | * | 12 - 17 Months | 102 | 13.7 |
| 18 - 23 Months | 19 | * | 18 - 23 Months | 29 | * |
| 24 - 35 Months | 38 | * | 24 - 35 Months | 91 | * |
| 36 - 59 Months | 99 | * | 36 - 59 Months | 334 | 22.2 |
| Total | 226 | 11.1 | Total | 595 | 21.0 |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 66 | * | 6 - 11 Months | 5 | * |
| 12 - 17 Months | 348 | 11.2 | 12 - 17 Months | 8 | * |
| 18 - 23 Months | 114 | 15.8 | 18 - 23 Months | 4 | * |
| 24 - 35 Months | 223 | 20.2 | 24 - 35 Months | 8 | * |
| 36 - 59 Months | 503 | 17.9 | 36 - 59 Months | 18 | * |
| Total | 1,254 | 16.4 | Total | 43 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 303 | 16.5 | 6 - 11 Months | 12 | * |
| 12 - 17 Months | 691 | 12.4 | 12 - 17 Months | 8 | * |
| 18 - 23 Months | 365 | 8.2 | 18 - 23 Months | 6 | * |
| 24 - 35 Months | 712 | 11.1 | 24 - 35 Months | 13 | * |
| 36 - 59 Months | 1,886 | 9.9 | 36 - 59 Months | 41 | * |
| Total | 3,957 | 10.9 | Total | 80 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 3 | * | 6 - 11 Months | 269 | 17.5 |
| 12 - 17 Months | 3 | * | 12 - 17 Months | 529 | 15.5 |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 200 | 14.5 |
| 24 - 35 Months | 2 | * | 24 - 35 Months | 404 | 12.6 |
| 36 - 59 Months | 7 | * | 36 - 59 Months | 772 | 10.2 |
| Total | 15 | * | Total | 2,174 | 13.2 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia I | ndicators |
|------------------------------|----------|---------------------------|
| | Number | Anemia Low Hb/Hct % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 718 | 18.4 |
| 12 - 17 Months | 1,738 | 13.1 |
| 18 - 23 Months | 737 | 11.3 |
| 24 - 35 Months | 1,491 | 13.7 |
| 36 - 59 Months | 3,660 | 12.1 |
| Total | 8,344 | 13.1 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age (3, | 4) | |
|--------------------------------|--------|---------------------------|-------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | l 1 | * |
| 12 - 17 Months | l i | * | 12 - 17 Months | Ιí | * |
| 18 - 23 Months | 1 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 3 | * | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 3 | * | 36 - 59 Months | 1 | * |
| Total | 8 | * | Total | 4 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 0 | * |
| Total | 1 | * | Total | 0 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 3 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 5 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 2 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 3 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 12 | * | 36 - 59 Months | 0 | * |
| Total | 25 | * | Total | 0 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 2 | * | 12 - 17 Months | 2 | * |
| 18 - 23 Months | 1 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 0 | * |
| Total | 3 | * | Total | 3 | * |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Run Date: 6/3/09 **Alpine**

Page:

| | Anemia I | ndicators |
|-----------------------------|----------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| otal All Race/Ethnic Groups | | |
| 6 - 11 Months | 5 | * |
| 12 - 17 Months | 11 | * |
| 18 - 23 Months | 4 | * |
| 24 - 35 Months | 8 | * |
| 36 - 59 Months | 16 | * |
| otal | 44 | * |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anomia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|---------------------------|----------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | by nace/Ellillicity and Age | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 18 | * | 6 - 11 Months | 1 1 | * |
| 12 - 17 Months | 30 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 4 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 17 | * | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 28 | * | 36 - 59 Months | 2 | * |
| Total | 97 | * | Total | 4 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | l 1 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 4 | * | 36 - 59 Months | 0 | * |
| Total | 5 | * | Total | 0 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 10 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 18 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 10 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 18 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 35 | * | 36 - 59 Months | 0 | * |
| Total | 91 | * | Total | 0 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 2 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 3 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 10 | * |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 6 | * |
| Total | 0 | * | Total | 22 | * |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age

Children Aged < 5 Years (2) County: 003

Page:

Run Date:

6/3/09

| | Anemia | Indicators | by Race | e/Ethnicity a | nd Age (3, 4 |) | |
|------------------------------|--------|----------------------|---------|---------------|--------------|--------|----------------------|
| | | Anemia Low Hb/Hct | | | | | Anemia Low Hb/Hct |
| | Number | % | | | | Number | % |
| Total All Race/Ethnic Groups | | | | | | | |
| 6 - 11 Months | 31 | * | | | | | |
| 12 - 17 Months | 52 | * | | | | | |
| 18 - 23 Months | 15 | * | | | | | |
| 24 - 35 Months | 46 | * | | | | | |
| 36 - 59 Months | 75 | * | | | | | |
| Total | 219 | 13.7 | | | | | |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4 |) | |
|--------------------------------|--------|------------|---------------------------------|--------|------------|
| | | Anemia | | | Anemia |
| | | Low Hb/Hct | | | Low Hb/Hct |
| | Number | % | | Number | % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 582 | 16.2 | 6 - 11 Months | 72 | * |
| 12 - 17 Months | 607 | 12.5 | 12 - 17 Months | 75 | * |
| 18 - 23 Months | 260 | 12.3 | 18 - 23 Months | 37 | * |
| 24 - 35 Months | 606 | 9.4 | 24 - 35 Months | 79 | * |
| 36 - 59 Months | 1,036 | 8.5 | 36 - 59 Months | 138 | 17.4 |
| Total | 3,091 | 11.2 | Total | 401 | 16.5 |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 29 | * | 6 - 11 Months | 3 | * |
| 12 - 17 Months | 38 | * | 12 - 17 Months | 4 | * |
| 18 - 23 Months | 17 | * | 18 - 23 Months | 2 | * |
| 24 - 35 Months | 35 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 73 | * | 36 - 59 Months | 5 | * |
| Total | 192 | 15.6 | Total | 14 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 124 | 21.8 | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 167 | 14.4 | 12 - 17 Months | 1 | * |
| 18 - 23 Months | 76 | * | 18 - 23 Months | 2 | * |
| 24 - 35 Months | 168 | 6.0 | 24 - 35 Months | 2 | * |
| 36 - 59 Months | 306 | 7.8 | 36 - 59 Months | 3 | * |
| Total | 841 | 11.1 | Total | 8 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 22 | * | 6 - 11 Months | 144 | 26.4 |
| 12 - 17 Months | 36 | * | 12 - 17 Months | 132 | 20.5 |
| 18 - 23 Months | 13 | * | 18 - 23 Months | 48 | * |
| 24 - 35 Months | 19 | * | 24 - 35 Months | 137 | 15.3 |
| 36 - 59 Months | 53 | * | 36 - 59 Months | 187 | 9.6 |
| Total | 143 | 10.5 | Total | 648 | 17.4 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

County: 004

Page:

Run Date:

6/3/09

| | Anemia I | ndicators |
|----------------------------------|----------------|---------------------------|
| | Number | Anemia Low Hb/Hct % |
| Total All Race/Ethnic Groups | | 10.5 |
| 6 - 11 Months 12 - 17 Months | 976 1.060 | 18.5 13.6 |
| 18 - 23 Months | 455 | 14.1 |
| 24 - 35 Months 36 - 59 Months | 1,046 1,801 | 10.8 9.2 |
| Total | 5,338 | 12.5 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| Anemia Indicators by Race/Ethnicity and Age (3, 4) | | | | | | | | | |
|--|--------|---------------------------|------------------|--------|---------------------------|--|--|--|--|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % | | | | |
| White, Not Hispanic | | | Asian | | | | | | |
| 6 - 11 Months | 23 | * | 6 - 11 Months | 1 | * | | | | |
| 12 - 17 Months | 37 | * | 12 - 17 Months | 4 | * | | | | |
| 18 - 23 Months | 12 | * | 18 - 23 Months | 1 | * | | | | |
| 24 - 35 Months | 22 | * | 24 - 35 Months | 4 | * | | | | |
| 36 - 59 Months | 39 | * | 36 - 59 Months | 8 | * | | | | |
| Total | 133 | 6.8 | Total | 18 | * | | | | |
| Black, Not Hispanic | | | Pacific Islander | | | | | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 0 | * | | | | |
| 12 - 17 Months | 1 | * | 12 - 17 Months | 0 | * | | | | |
| 18 - 23 Months | 1 | * | 18 - 23 Months | 0 | * | | | | |
| 24 - 35 Months | 2 | * | 24 - 35 Months | 0 | * | | | | |
| 36 - 59 Months | 5 | * | 36 - 59 Months | 1 | * | | | | |
| Total | 10 | * | Total | 1 | * | | | | |
| Hispanic | | | Filipino | | | | | | |
| 6 - 11 Months | 75 | * | 6 - 11 Months | 0 | * | | | | |
| 12 - 17 Months | 71 | * | 12 - 17 Months | 0 | * | | | | |
| 18 - 23 Months | 35 | * | 18 - 23 Months | 0 | * | | | | |
| 24 - 35 Months | 73 | * | 24 - 35 Months | 2 | * | | | | |
| 36 - 59 Months | 172 | 9.9 | 36 - 59 Months | 3 | * | | | | |
| Total | 426 | 14.3 | Total | 5 | * | | | | |
| American Indian/Alaskan Native | | | All Other | | | | | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 53 | * | | | | |
| 12 - 17 Months | 1 | * | 12 - 17 Months | 45 | * | | | | |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 30 | * | | | | |
| 24 - 35 Months | 1 | * | 24 - 35 Months | 36 | * | | | | |
| 36 - 59 Months | 1 | * | 36 - 59 Months | 104 | 12.5 | | | | |
| Total | 3 | * | Total | 268 | 13.1 | | | | |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Children Aged < 5 Years (2)

Anemia Indicators by Race/Ethnicity and Age

Run Date: 6/3/09

Page:

| | Anemia I | ndicators | 1) |
|-----------------------------|----------|----------------------|--------|
| | | Anemia Low Hb/Hct | |
| | Number | % | Number |
| otal All Race/Ethnic Groups | | | • |
| 6 - 11 Months | 153 | 15.0 | |
| 12 - 17 Months | 159 | 13.8 | |
| 18 - 23 Months | 79 | * | |
| 24 - 35 Months | 140 | 8.6 | |
| 36 - 59 Months | 333 | 10.5 | |
| otal | 864 | 12.3 | |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 17B 2008 Pediatric Nutrition Surveillance (1) City Of Berkeley

Page:

Run Date:

6/3/09

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | Anemia | Indicators | y Race/Ethnicity and Age (3, | 4) | |
|--------------------------------|--------|---------------------------|------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 1 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 2 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 0 | * |
| Total | 3 | * | Total | 0 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 1 | * | 36 - 59 Months | 0 | * |
| Total | 1 | * | Total | 0 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 5 | * | 6 - 11 Months | 1 | * |
| 12 - 17 Months | 5 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 2 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 3 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 6 | * | 36 - 59 Months | 0 | * |
| Total | 21 | * | Total | 1 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 4 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 2 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 6 | * |
| Total | 0 | * | Total | 13 | * |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 17B

2008 Pediatric Nutrition Surveillance (1) City Of Berkeley

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

County: 059

| | Anemia II | ndicators | (3, 4) | |
|------------------------------|-----------|----------------------|--------|---------|
| | | Anemia Low Hb/Hct | | A Lo |
| | Number | % | Number | r |
| Total All Race/Ethnic Groups | | | | |
| 6 - 11 Months | 10 | * | | |
| 12 - 17 Months | 8 | * | | |
| 18 - 23 Months | 4 | * | | |
| 24 - 35 Months | 4 | * | | |
| 36 - 59 Months | 13 | * | | |
| Total | 39 | * | | |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|---------------------------|----------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 9 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 10 | * | 12 - 17 Months | 1 | * |
| 18 - 23 Months | 5 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 7 | * | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 15 | * | 36 - 59 Months | 1 | * |
| Total | 46 | * | Total | 3 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 1 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 1 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 1 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 8 | * | 36 - 59 Months | 0 | * |
| Total | 12 | * | Total | 0 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 146 | 30.1 | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 127 | 22.8 | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 44 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 54 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 184 | 14.7 | 36 - 59 Months | 1 | * |
| Total | 555 | 21.4 | Total | 1 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 10 | * |
| 12 - 17 Months | 1 | * | 12 - 17 Months | 14 | * |
| 18 - 23 Months | 1 | * | 18 - 23 Months | 2 | * |
| 24 - 35 Months | 1 | * | 24 - 35 Months | 19 | * |
| 36 - 59 Months | 2 | * | 36 - 59 Months | 33 | * |
| Total | 5 | * | Total | 78 | * |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

County: 006

Page:

Run Date:

6/3/09

| | Anemia I | ndicators |
|-----------------------------|----------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| otal All Race/Ethnic Groups | | |
| 6 - 11 Months | 166 | 31.9 |
| 12 - 17 Months | 154 | 22.1 |
| 18 - 23 Months | 53 | * |
| 24 - 35 Months | 83 | * |
| 36 - 59 Months | 244 | 15.6 |
| Total | 700 | 21.9 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 17B 2008 Pediatric Nutrition Surveillance (1) Contra Costa

Page:

Run Date:

6/3/09

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|-----------|---------------------------|----------------------------------|----------|---------------------------|
| | Neuraleau | Anemia Low Hb/Hct % | | Niversia | Anemia Low Hb/Hct % |
| | Number | /0 | | Number | /0 |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 46 | * | 6 - 11 Months | 7 | * |
| 12 - 17 Months | 40 | * | 12 - 17 Months | 6 | * |
| 18 - 23 Months | 18 | * | 18 - 23 Months | 2 | * |
| 24 - 35 Months | 28 | * | 24 - 35 Months | 2 | * |
| 36 - 59 Months | 74 | * | 36 - 59 Months | 19 | * |
| Total | 206 | 16.0 | Total | 36 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 56 | * | 6 - 11 Months | 5 | * |
| 12 - 17 Months | 86 | * | 12 - 17 Months | 2 | * |
| 18 - 23 Months | 20 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 71 | * | 24 - 35 Months | 4 | * |
| 36 - 59 Months | 163 | 21.5 | 36 - 59 Months | 4 | * |
| Total | 396 | 24.2 | Total | 15 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 294 | 26.9 | 6 - 11 Months | 6 | * |
| 12 - 17 Months | 468 | 16.0 | 12 - 17 Months | 4 | * |
| 18 - 23 Months | 117 | 20.5 | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 317 | 11.4 | 24 - 35 Months | 8 | * |
| 36 - 59 Months | 677 | 11.8 | 36 - 59 Months | 16 | * |
| Total | 1,873 | 15.7 | Total | 34 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 192 | 13.5 |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 93 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 40 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 123 | 15.4 |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 274 | 12.8 |
| Total | 1 | * | Total | 722 | 13.3 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 17B

2008 Pediatric Nutrition Surveillance (1) Contra Costa

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

County: 007

Page:

Run Date:

6/3/09

| | Anemia I | ndicators |
|----------------------------------|------------|---------------------------|
| | Number | Anemia Low Hb/Hct % |
| Total All Race/Ethnic Groups | Number | /0 |
| 6 - 11 Months | 607 | 21.9 |
| 12 - 17 Months 18 - 23 Months | 699 197 | 16.7 17.8 |
| 24 - 35 Months | 553 | 15.2 |
| 36 - 59 Months | 1,227 | 13.2 |
| Total | 3,283 | 16.2 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page:

Run Date:

6/3/09

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | Anemia l | ndicators | y Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|----------|---------------------------|---------------------------------|--------|---------------------------|
| | | Anemia Low Hb/Hct % | | | Anemia Low Hb/Hct % |
| | Number | 70 | | Number | 70 |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 21 | * | 6 - 11 Months | 1 | * |
| 12 - 17 Months | 69 | * | 12 - 17 Months | 10 | * |
| 18 - 23 Months | 23 | * | 18 - 23 Months | 6 | * |
| 24 - 35 Months | 55 | * | 24 - 35 Months | 9 | * |
| 36 - 59 Months | 86 | * | 36 - 59 Months | 20 | * |
| Total | 254 | 5.5 | Total | 46 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 3 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 1 | * | 36 - 59 Months | 1 | * |
| Total | 4 | * | Total | 1 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 11 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 28 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 9 | * | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 19 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 39 | * | 36 - 59 Months | 1 | * |
| Total | 106 | 5.7 | Total | 2 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 4 | * | 6 - 11 Months | 6 | * |
| 12 - 17 Months | 15 | * | 12 - 17 Months | 7 | * |
| 18 - 23 Months | 6 | * | 18 - 23 Months | 6 | * |
| 24 - 35 Months | 13 | * | 24 - 35 Months | 14 | * |
| 36 - 59 Months | 24 | * | 36 - 59 Months | 31 | * |
| Total | 62 | * | Total | 64 | * |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page:

Run Date:

6/3/09

Del Norte

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | Anemia I | ndicators |
|------------------------------|----------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 43 | * |
| 12 - 17 Months | 132 | 8.3 |
| 18 - 23 Months | 51 | * |
| 24 - 35 Months | 110 | 10.0 |
| 36 - 59 Months | 203 | 4.9 |
| Total | 539 | 6.7 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

State: California County: 009

Page:

Run Date:

6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4 | 4) | |
|--------------------------------|--------|---------------------------|---------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 88 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 73 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 33 | * | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 98 | * | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 187 | 12.3 | 36 - 59 Months | 2 | * |
| Total | 479 | 14.0 | Total | 4 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 3 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 1 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 1 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 3 | * | 36 - 59 Months | 0 | * |
| Total | 9 | * | Total | 0 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 127 | 15.0 | 6 - 11 Months | 1 | * |
| 12 - 17 Months | 115 | 13.9 | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 29 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 127 | 19.7 | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 197 | 8.1 | 36 - 59 Months | 1 | * |
| Total | 595 | 13.4 | Total | 3 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 71 | * |
| 12 - 17 Months | 1 | * | 12 - 17 Months | 66 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 17 | * |
| 24 - 35 Months | 1 | * | 24 - 35 Months | 45 | * |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 77 | * |
| Total | 3 | * | Total | 276 | 12.0 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

El Dorado Anemia Indicators by Race/Ethnicity and Age Page: Run Date: 6/3/09

Children Aged < 5 Years (2)

| | Anemia I | Indicators |
|------------------------------|----------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 289 | 13.5 |
| 12 - 17 Months | 258 | 14.3 |
| 18 - 23 Months | 81 | * |
| 24 - 35 Months | 274 | 16.4 |
| 36 - 59 Months | 467 | 10.1 |
| Total | 1,369 | 13.3 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age

Children Aged < 5 Years (2) County: 010

Page:

Run Date:

6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age $^{(3,4)}$ | | |
|----------------------------------|--------|---------------------------|--------------------------------------|----------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | Number | | Asian | INGINIDO | |
| 6 - 11 Months | 714 | 12.7 | 6 - 11 Months | 215 | 20.5 |
| 12 - 17 Months | 706 | 11.8 | 12 - 17 Months | 392 | 20.3 17.9 |
| 18 - 23 Months | 302 | 9.3 | 18 - 23 Months | 136 | 17.9 |
| 24 - 35 Months | 576 | 13.5 | 24 - 35 Months | 361 | 32.7 |
| 36 - 59 Months | 1,131 | 7.2 | 36 - 59 Months | 678 | 19.0 |
| Total | 3,429 | 10.5 | Total | 1,782 | 21.7 |
| | 3,429 | 10.5 | Pacific Islander | 1,702 | 21.1 |
| Black, Not Hispanic | 245 | 18.8 | | 3 | * |
| 6 - 11 Months | 281 | 14.2 | 6 - 11 Months | 5 | * |
| 12 - 17 Months 18 - 23 Months | 166 | 15.1 | 12 - 17 Months 18 - 23 Months | 4 | * |
| 24 - 35 Months | 267 | 19.5 | 24 - 35 Months | 4 | * |
| 36 - 59 Months | 479 | 16.3 | 36 - 59 Months | 4 | * |
| Total | 1,438 | 16.8 | Total | 20 | * |
| | 1,430 | 10.0 | | 20 | |
| Hispanic 6 - 11 Months | 4.680 | 22.2 | Filipino 6 - 11 Months | 9 | * |
| 12 - 17 Months | 5,321 | 13.5 | 12 - 17 Months | 3 | * |
| 18 - 23 Months | 2,484 | 14.5 | 12 - 17 Months 18 - 23 Months | 2 | * |
| 24 - 35 Months | 4,565 | 15.0 | 24 - 35 Months | 6 | * |
| 36 - 59 Months | 8,830 | 10.1 | 36 - 59 Months | 4 | * |
| Total | 25,880 | 14.2 | Total | 24 | * |
| American Indian/Alaskan Native | 23,000 | 17.2 | All Other | 27 | |
| 6 - 11 Months | 38 | * | 6 - 11 Months | 721 | 24.7 |
| 12 - 17 Months | 62 | * | 12 - 17 Months | 622 | 15.1 |
| 18 - 23 Months | 17 | * | 18 - 23 Months | 246 | 19.1 |
| 24 - 35 Months | 45 | * | 24 - 35 Months | 556 | 14.7 |
| 36 - 59 Months | 82 | * | 36 - 59 Months | 1,202 | 13.9 |
| Total | 244 | 9.4 | Total | 3,347 | 17.0 |
| IUlai | 244 | 9.4 | IUlai | 3,347 | 17.0 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators |
|------------------------------|--------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 6,625 | 21.2 |
| 12 - 17 Months | 7,392 | 13.7 |
| 18 - 23 Months | 3,357 | 14.5 |
| 24 - 35 Months | 6,380 | 16.0 |
| 36 - 59 Months | 12,410 | 10.9 |
| Total | 36,164 | 14.6 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page:

Run Date:

6/3/09

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | | | (0.4) | | |
|--------------------------------|--------|---------------------------|----------------------------------|--------|---------------------------|
| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White Net Hieronia | Number | ,,, | Asian | Number | |
| White, Not Hispanic | 81 | * | Asian | _ | * |
| 6 - 11 Months | 70 | * | 6 - 11 Months | 5 2 | * |
| 12 - 17 Months | 36 | * | 12 - 17 Months | 2 | * |
| 18 - 23 Months | 65 | * | 18 - 23 Months | 6 | * |
| 24 - 35 Months | 102 | 0.0 | 24 - 35 Months | 10 | * |
| 36 - 59 Months | | 9.8 | 36 - 59 Months | _ | |
| Total | 354 | 18.6 | Total | 25 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 2 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 2 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 2 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 2 | * | 36 - 59 Months | 0 | * |
| Total | 8 | * | Total | 0 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 115 | 27.0 | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 109 | 22.9 | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 70 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 103 | 22.3 | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 204 | 8.8 | 36 - 59 Months | 0 | * |
| Total | 601 | 18.0 | Total | 0 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 45 | * |
| 12 - 17 Months | 3 | * | 12 - 17 Months | 25 | * |
| 18 - 23 Months | 3 | * | 18 - 23 Months | 8 | * |
| 24 - 35 Months | 2 | * | 24 - 35 Months | 33 | * |
| 36 - 59 Months | 6 | * | 36 - 59 Months | 36 | * |
| Total | 15 | * | Total | 147 | 19.7 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia I | ndicators |
|------------------------------|----------|---------------------------|
| | Number | Anemia Low Hb/Hct % |
| Total All Race/Ethnic Groups | Number | /6 |
| 6 - 11 Months | 249 | 28.1 |
| 12 - 17 Months | 211 | 22.3 |
| 18 - 23 Months | 119 | 16.8 |
| 24 - 35 Months | 211 | 21.8 |
| 36 - 59 Months | 360 | 9.7 |
| Total | 1,150 | 19.0 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

County: 012

Page:

Run Date:

6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|----------------------|----------------------------------|--------|----------------------|
| | | Anemia Low Hb/Hct | | | Anemia Low Hb/Hct |
| | Number | % | | Number | % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 169 | 21.9 | 6 - 11 Months | 6 | * |
| 12 - 17 Months | 125 | 18.4 | 12 - 17 Months | 7 | * |
| 18 - 23 Months | 72 | * | 18 - 23 Months | 5 | * |
| 24 - 35 Months | 117 | 26.5 | 24 - 35 Months | 6 | * |
| 36 - 59 Months | 198 | 13.1 | 36 - 59 Months | 8 | * |
| Total | 681 | 18.2 | Total | 32 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 3 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 1 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 4 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 1 | * | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 4 | * | 36 - 59 Months | 1 | * |
| Total | 13 | * | Total | 2 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 48 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 39 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 29 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 54 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 85 | * | 36 - 59 Months | 0 | * |
| Total | 255 | 10.2 | Total | 0 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 14 | * | 6 - 11 Months | 223 | 9.4 |
| 12 - 17 Months | 37 | * | 12 - 17 Months | 209 | 5.3 |
| 18 - 23 Months | 11 | * | 18 - 23 Months | 78 | * |
| 24 - 35 Months | 32 | * | 24 - 35 Months | 187 | 8.0 |
| 36 - 59 Months | 78 | * | 36 - 59 Months | 277 | 5.4 |
| Total | 172 | 11.0 | Total | 974 | 6.8 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Humboldt

Page:

Run Date:

6/3/09

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | Anemia I | ndicators |
|----------------------------------|------------|---------------------------|
| | Neuralaan | Anemia Low Hb/Hct % |
| al All Race/Ethnic Groups | Number | 70 |
| 6 - 11 Months | 463 | 14.9 |
| 12 - 17 Months | 418 199 | 10.3 |
| 18 - 23 Months 24 - 35 Months | 398 | 7.5 14.3 |
| 36 - 59 Months | 651 | 8.8 |
| Total | 2,129 | 11.3 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators | y Race/Ethnicity and Age (3, 4 |) | |
|--------------------------------|--------|----------------------|--------------------------------|--------|----------------------|
| | | Anemia Low Hb/Hct | | | Anemia Low Hb/Hct |
| | Number | % | | Number | % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 14 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 15 | * | 12 - 17 Months | 3 | * |
| 18 - 23 Months | 8 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 15 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 48 | * | 36 - 59 Months | 1 | * |
| Total | 100 | 15.0 | Total | 4 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 3 | * | 6 - 11 Months | 2 | * |
| 12 - 17 Months | 9 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 2 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 12 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 25 | * | 36 - 59 Months | 1 | * |
| Total | 51 | * | Total | 3 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 686 | 8.9 | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 1,110 | 5.8 | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 393 | 8.4 | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 1,096 | 7.8 | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 2,176 | 5.7 | 36 - 59 Months | 2 | * |
| Total | 5,461 | 6.7 | Total | 2 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 22 | * |
| 12 - 17 Months | 1 | * | 12 - 17 Months | 39 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 13 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 23 | * |
| 36 - 59 Months | 1 | * | 36 - 59 Months | 52 | * |
| Total | 2 | * | Total | 149 | 10.1 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age

Children Aged < 5 Years (2) County: 013

Page:

Run Date:

6/3/09

| | Anemia In | ndicators |
|------------------------------|-----------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 727 | 9.4 |
| 12 - 17 Months | 1,177 | 6.2 |
| 18 - 23 Months | 416 | 8.7 |
| 24 - 35 Months | 1,146 | 8.0 |
| 36 - 59 Months | 2,306 | 5.9 |
| Total | 5,772 | 7.0 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age (3, | 4) | |
|-------------------------------|--------|---------------------------|-------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 3 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 4 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 2 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 3 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 7 | * | 36 - 59 Months | 0 | * |
| Total | 19 | * | Total | 0 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 2 | * | 36 - 59 Months | 0 | * |
| Total | 2 | * | Total | 0 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 5 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 19 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 14 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 20 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 21 | * | 36 - 59 Months | 0 | * |
| Total | 79 | * | Total | 0 | * |
| American Indian/Alaskan Nativ | 'e | | All Other | | |
| 6 - 11 Months | 7 | * | 6 - 11 Months | 2 | * |
| 12 - 17 Months | 6 | * | 12 - 17 Months | 1 | * |
| 18 - 23 Months | 1 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 1 | * | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 9 | * | 36 - 59 Months | 3 | * |
| Total | 24 | * | Total | 7 | * |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page:

Run Date:

6/3/09

Inyo

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | Anemia I | ndicators |
|----------------------------------|----------|---------------------------|
| | Number | Anemia Low Hb/Hct % |
| Total All Race/Ethnic Groups | Number | |
| 6 - 11 Months 12 - 17 Months | 17 30 | * |
| 18 - 23 Months | 17 | * |
| 24 - 35 Months 36 - 59 Months | 25 42 | * |
| Total | 131 | 12.2 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page:

Run Date:

6/3/09

County: 015

| | Anemia | Indicators | y Race/Ethnicity and Age (3, 4 |) | |
|--------------------------------|--------|------------|--------------------------------|--------|------------|
| | | Anemia | | | Anemia |
| | | Low Hb/Hct | | | Low Hb/Hct |
| | Number | % | | Number | % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 279 | 15.8 | 6 - 11 Months | 10 | * |
| 12 - 17 Months | 311 | 11.9 | 12 - 17 Months | 14 | * |
| 18 - 23 Months | 146 | 16.4 | 18 - 23 Months | 3 | * |
| 24 - 35 Months | 336 | 10.7 | 24 - 35 Months | 15 | * |
| 36 - 59 Months | 632 | 8.7 | 36 - 59 Months | 28 | * |
| Total | 1,704 | 11.5 | Total | 70 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 131 | 21.4 | 6 - 11 Months | 2 | * |
| 12 - 17 Months | 170 | 21.2 | 12 - 17 Months | 1 | * |
| 18 - 23 Months | 79 | * | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 188 | 31.9 | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 362 | 19.6 | 36 - 59 Months | 6 | * |
| Total | 930 | 23.1 | Total | 11 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 2,191 | 17.1 | 6 - 11 Months | 8 | * |
| 12 - 17 Months | 2,618 | 12.4 | 12 - 17 Months | 6 | * |
| 18 - 23 Months | 1,159 | 13.2 | 18 - 23 Months | 4 | * |
| 24 - 35 Months | 2,793 | 13.0 | 24 - 35 Months | 11 | * |
| 36 - 59 Months | 5,679 | 9.1 | 36 - 59 Months | 25 | * |
| Total | 14,440 | 12.0 | Total | 54 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 185 | 19.5 |
| 12 - 17 Months | 2 | * | 12 - 17 Months | 201 | 17.9 |
| 18 - 23 Months | 1 | * | 18 - 23 Months | 99 | * |
| 24 - 35 Months | 2 | * | 24 - 35 Months | 211 | 17.1 |
| 36 - 59 Months | 5 | * | 36 - 59 Months | 453 | 10.2 |
| Total | 11 | * | Total | 1,149 | 14.7 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Kern Anemia Indicators by Race/Ethnicity and Age

Children Aged < 5 Years (2) County: 015

Page:

Run Date:

6/3/09

| | Anemia I | ndicators | y Race/Ethnici | ity and Age (3, 4) | | |
|------------------------------|----------|---------------------------|----------------|--------------------|--------|--------------|
| | Number | Anemia Low Hb/Hct % | | | Number | Ane Low F |
| Total All Race/Ethnic Groups | | | | | | |
| 6 - 11 Months | 2,807 | 17.3 | | | | |
| 12 - 17 Months | 3,323 | 13.2 | | | | |
| 18 - 23 Months | 1,492 | 14.2 | | | | |
| 24 - 35 Months | 3,557 | 14.0 | | | | |
| 36 - 59 Months | 7,190 | 9.7 | | | | |
| Total | 18,369 | 12.7 | | | | |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|---------------------------|----------------------------------|---------|---------------------------|
| | | Anemia Low Hb/Hct % | | <u></u> | Anemia Low Hb/Hct % |
| | Number | 70 | | Number | 76 |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 43 | * | 6 - 11 Months | 4 | * |
| 12 - 17 Months | 57 | * | 12 - 17 Months | 5 | * |
| 18 - 23 Months | 33 | * | 18 - 23 Months | 2 | * |
| 24 - 35 Months | 55 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 84 | * | 36 - 59 Months | 1 | * |
| Total | 272 | 25.0 | Total | 12 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 19 | * | 6 - 11 Months | 1 | * |
| 12 - 17 Months | 20 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 4 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 29 | * | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 36 | * | 36 - 59 Months | 3 | * |
| Total | 108 | 36.1 | Total | 5 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 426 | 32.6 | 6 - 11 Months | 1 | * |
| 12 - 17 Months | 537 | 29.2 | 12 - 17 Months | 2 | * |
| 18 - 23 Months | 224 | 26.8 | 18 - 23 Months | 2 | * |
| 24 - 35 Months | 412 | 25.5 | 24 - 35 Months | 2 | * |
| 36 - 59 Months | 760 | 19.7 | 36 - 59 Months | 4 | * |
| Total | 2,359 | 25.9 | Total | 11 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 55 | * |
| 12 - 17 Months | 2 | * | 12 - 17 Months | 102 | 27.5 |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 39 | * |
| 24 - 35 Months | 6 | * | 24 - 35 Months | 89 | * |
| 36 - 59 Months | 2 | * | 36 - 59 Months | 150 | 15.3 |
| Total | 11 | * | Total | 435 | 22.5 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia I | ndicators |
|------------------------------|----------|---------------------------|
| | Number | Anemia Low Hb/Hct % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 550 | 32.7 |
| 12 - 17 Months | 725 | 28.1 |
| 18 - 23 Months | 304 | 27.6 |
| 24 - 35 Months | 594 | 26.1 |
| 36 - 59 Months | 1,040 | 19.3 |
| Total | 3,213 | 25.6 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page:

Run Date:

6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|-----------|---------------------------|----------------------------------|-----------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | 110111001 | | Asian | - Trambon | |
| 6 - 11 Months | 44 | * | 6 - 11 Months | 35 | * |
| 12 - 17 Months | 37 | * | 12 - 17 Months | 42 | * |
| 18 - 23 Months | 12 | * | 18 - 23 Months | 10 | * |
| 24 - 35 Months | 26 | * | 24 - 35 Months | 31 | * |
| 36 - 59 Months | 61 | * | 36 - 59 Months | 96 | * |
| Total | 180 | 16.7 | Total | 214 | 8.9 |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 39 | * | 6 - 11 Months | 2 | * |
| 12 - 17 Months | 19 | * | 12 - 17 Months | 1 | * |
| 18 - 23 Months | 11 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 24 | * | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 58 | * | 36 - 59 Months | 2 | * |
| Total | 151 | 15.9 | Total | 6 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 2,029 | 22.9 | 6 - 11 Months | 7 | * |
| 12 - 17 Months | 2,474 | 16.2 | 12 - 17 Months | 2 | * |
| 18 - 23 Months | 1,071 | 17.1 | 18 - 23 Months | 2 | * |
| 24 - 35 Months | 2,729 | 17.0 | 24 - 35 Months | 6 | * |
| 36 - 59 Months | 5,586 | 12.1 | 36 - 59 Months | 20 | * |
| Total | 13,889 | 15.7 | Total | 37 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 89 | * |
| 12 - 17 Months | 1 | * | 12 - 17 Months | 116 | 17.2 |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 28 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 105 | 17.1 |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 228 | 12.3 |
| Total | 1 | * | Total | 566 | 16.4 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators |
|--------------------------------|------------------------|---------------------------|
| | Number | Anemia Low Hb/Hct % |
| Total All Race/Ethnic Groups | Number | /6 |
| 6 - 11 Months | 2,245 | 22.2 |
| 12 - 17 Months | 2,692 | 16.2 |
| 18 - 23 Months | 1,134 | 16.9 |
| 24 - 35 Months | 2,922 | 16.8 |
| 36 - 59 Months Total | 6,051 15,044 | 12.2 15.6 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

County: 094

Page:

Run Date:

6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|---------------------------|----------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| | Number | /0 | | Number | /0 |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 69 | * | 6 - 11 Months | 3 | * |
| 12 - 17 Months | 43 | * | 12 - 17 Months | 5 | * |
| 18 - 23 Months | 23 | * | 18 - 23 Months | 2 | * |
| 24 - 35 Months | 57 | * | 24 - 35 Months | 2 | * |
| 36 - 59 Months | 134 | 13.4 | 36 - 59 Months | 14 | * |
| Total | 326 | 14.1 | Total | 26 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 18 | * | 6 - 11 Months | 2 | * |
| 12 - 17 Months | 13 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 10 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 26 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 55 | * | 36 - 59 Months | 3 | * |
| Total | 122 | 18.9 | Total | 5 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 635 | 17.3 | 6 - 11 Months | 6 | * |
| 12 - 17 Months | 933 | 11.8 | 12 - 17 Months | 1 | * |
| 18 - 23 Months | 418 | 12.4 | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 1,026 | 13.8 | 24 - 35 Months | 7 | * |
| 36 - 59 Months | 2,272 | 9.0 | 36 - 59 Months | 7 | * |
| Total | 5,284 | 11.7 | Total | 22 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 86 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 71 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 16 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 54 | * |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 146 | 13.0 |
| Total | 0 | * | Total | 373 | 16.6 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page:

Run Date:

6/3/09

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | Anemia I | ndicators | |
|-----------------------------|----------|----------------------|--------|
| | | Anemia Low Hb/Hct | |
| | Number | % | Number |
| otal All Race/Ethnic Groups | | | • |
| 6 - 11 Months | 819 | 18.4 | |
| 12 - 17 Months | 1,066 | 12.2 | |
| 18 - 23 Months | 470 | 12.3 | |
| 24 - 35 Months | 1,172 | 13.8 | |
| 36 - 59 Months | 2,631 | 9.7 | |
| otal | 6,158 | 12.3 | |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

County: 097

Page:

Run Date:

6/3/09

| | Anemia | Indicators | y Race/Ethnicity and Age (3, 4 | 4) | |
|--------------------------------|----------|---------------------------|--------------------------------|----------|---------------------------|
| | Niverban | Anemia Low Hb/Hct % | | Newsland | Anemia Low Hb/Hct % |
| | Number | /0 | | Number | /0 |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 397 | 14.6 | 6 - 11 Months | 218 | 15.6 |
| 12 - 17 Months | 345 | 14.5 | 12 - 17 Months | 222 | 6.8 |
| 18 - 23 Months | 146 | 9.6 | 18 - 23 Months | 77 | * |
| 24 - 35 Months | 348 | 13.2 | 24 - 35 Months | 225 | 10.2 |
| 36 - 59 Months | 676 | 7.8 | 36 - 59 Months | 467 | 9.2 |
| Total | 1,912 | 11.6 | Total | 1,209 | 10.2 |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 486 | 18.9 | 6 - 11 Months | 14 | * |
| 12 - 17 Months | 758 | 21.1 | 12 - 17 Months | 14 | * |
| 18 - 23 Months | 330 | 19.1 | 18 - 23 Months | 12 | * |
| 24 - 35 Months | 797 | 20.6 | 24 - 35 Months | 19 | * |
| 36 - 59 Months | 1,529 | 18.5 | 36 - 59 Months | 47 | * |
| Total | 3,900 | 19.5 | Total | 106 | 10.4 |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 10,981 | 19.2 | 6 - 11 Months | 53 | * |
| 12 - 17 Months | 15,206 | 14.4 | 12 - 17 Months | 64 | * |
| 18 - 23 Months | 5,933 | 12.5 | 18 - 23 Months | 25 | * |
| 24 - 35 Months | 14,542 | 13.8 | 24 - 35 Months | 60 | * |
| 36 - 59 Months | 27,869 | 9.2 | 36 - 59 Months | 131 | 11.5 |
| Total | 74,531 | 12.9 | Total | 333 | 13.2 |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 6 | * | 6 - 11 Months | 928 | 18.3 |
| 12 - 17 Months | 22 | * | 12 - 17 Months | 1,107 | 15.4 |
| 18 - 23 Months | 10 | * | 18 - 23 Months | 430 | 17.4 |
| 24 - 35 Months | 15 | * | 24 - 35 Months | 1,095 | 17.4 |
| 36 - 59 Months | 33 | * | 36 - 59 Months | 2,018 | 12.0 |
| Total | 86 | * | Total | 5,578 | 15.2 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page:

Run Date:

6/3/09

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | Anemia I | ndicators |
|------------------------------|----------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 13,083 | 18.9 |
| 12 - 17 Months | 17,738 | 14.7 |
| 18 - 23 Months | 6,963 | 13.1 |
| 24 - 35 Months | 17,101 | 14.3 |
| 36 - 59 Months | 32,770 | 9.8 |
| Total | 87,655 | 13.3 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

State: California County: 096

Page:

Run Date:

6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|---------------------------|----------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 31 | * | 6 - 11 Months | 12 | * |
| 12 - 17 Months | 20 | * | 12 - 17 Months | 9 | * |
| 18 - 23 Months | 10 | * | 18 - 23 Months | 5 | * |
| 24 - 35 Months | 25 | * | 24 - 35 Months | 17 | * |
| 36 - 59 Months | 63 | * | 36 - 59 Months | 35 | * |
| Total | 149 | 10.7 | Total | 78 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 137 | 24.8 | 6 - 11 Months | 4 | * |
| 12 - 17 Months | 166 | 22.3 | 12 - 17 Months | 2 | * |
| 18 - 23 Months | 62 | * | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 176 | 21.0 | 24 - 35 Months | 6 | * |
| 36 - 59 Months | 355 | 17.7 | 36 - 59 Months | 13 | * |
| Total | 896 | 20.8 | Total | 26 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 1,935 | 18.9 | 6 - 11 Months | 11 | * |
| 12 - 17 Months | 2,076 | 16.2 | 12 - 17 Months | 12 | * |
| 18 - 23 Months | 1,060 | 14.1 | 18 - 23 Months | 3 | * |
| 24 - 35 Months | 2,405 | 14.2 | 24 - 35 Months | 18 | * |
| 36 - 59 Months | 4,931 | 9.1 | 36 - 59 Months | 33 | * |
| Total | 12,407 | 13.2 | Total | 77 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 85 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 96 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 42 | * |
| 24 - 35 Months | 2 | * | 24 - 35 Months | 113 | 15.9 |
| 36 - 59 Months | 4 | * | 36 - 59 Months | 312 | 9.0 |
| Total | 7 | * | Total | 648 | 12.8 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

L.A. South

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

County: 096

Page:

Run Date:

6/3/09

| Anemia Indicators by Race/Ethnicity and Age (3, 4) | | | | | | |
|--|--------|----------------------|--|--|--|--|
| | | Anemia Low Hb/Hct | | | | |
| | Number | % | | | | |
| Total All Race/Ethnic Groups | | | | | | |
| 6 - 11 Months | 2,216 | 19.1 | | | | |
| 12 - 17 Months | 2,381 | 16.8 | | | | |
| 18 - 23 Months | 1,183 | 14.3 | | | | |
| 24 - 35 Months | 2,762 | 14.7 | | | | |
| 36 - 59 Months | 5,746 | 9.6 | | | | |
| Total | 14,288 | 13.6 | | | | |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page:

Run Date:

6/3/09

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | Anemia | Indicators | by Race/Ethnicity and Age (3, | 4) | |
|--------------------------------|--------|---------------------------|-------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 10 | * | 6 - 11 Months | 1 | * |
| 12 - 17 Months | 4 | * | 12 - 17 Months | 3 | * |
| 18 - 23 Months | 6 | * | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 10 | * | 24 - 35 Months | 3 | * |
| 36 - 59 Months | 24 | * | 36 - 59 Months | 8 | * |
| Total | 54 | * | Total | 16 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 36 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 33 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 29 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 52 | * | 24 - 35 Months | 2 | * |
| 36 - 59 Months | 140 | 10.7 | 36 - 59 Months | 3 | * |
| Total | 290 | 15.2 | Total | 5 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 130 | 21.5 | 6 - 11 Months | 2 | * |
| 12 - 17 Months | 235 | 14.5 | 12 - 17 Months | l 1 | * |
| 18 - 23 Months | 86 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 225 | 11.6 | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 489 | 6.3 | 36 - 59 Months | 4 | * |
| Total | 1,165 | 11.0 | Total | 7 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 9 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 11 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 6 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 13 | * |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 42 | * |
| Total | 1 | * | Total | 81 | * |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page:

Run Date:

6/3/09

| | Anemia l | ndicators |
|-----------------------------|----------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| otal All Race/Ethnic Groups | | |
| 6 - 11 Months | 189 | 21.2 |
| 12 - 17 Months | 287 | 17.1 |
| 18 - 23 Months | 128 | 12.5 |
| 24 - 35 Months | 305 | 13.8 |
| 36 - 59 Months | 710 | 7.5 |
| otal | 1,619 | 12.4 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|---------------------------|----------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| AA/L': NI LIP | Number | | A : | Number | |
| White, Not Hispanic | 00 | * | Asian | | * |
| 6 - 11 Months | 39 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 64 | * | 12 - 17 Months | 0 | |
| 18 - 23 Months | 25 | * | 18 - 23 Months | 0 | |
| 24 - 35 Months | 34 | | 24 - 35 Months | 0 | |
| 36 - 59 Months | 79 | | 36 - 59 Months | ' | |
| Total | 241 | 10.0 | Total | 1 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 1 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 3 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 3 | * | 36 - 59 Months | 1 | * |
| Total | 7 | * | Total | 2 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 19 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 27 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 11 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 28 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 68 | * | 36 - 59 Months | 0 | * |
| Total | 153 | 7.8 | Total | 0 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 38 | * |
| 12 - 17 Months | 2 | * | 12 - 17 Months | 31 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 13 | * |
| 24 - 35 Months | 1 | * | 24 - 35 Months | 28 | * |
| 36 - 59 Months | 1 | * | 36 - 59 Months | 32 | * |
| Total | 4 | * | Total | 142 | 25.4 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Lake Anemia Indicators by Race/Ethnicity and Age Page: Run Date: 6/3/09

County: 017

Children Aged < 5 Years (2)

| Anemia Indicators by Race/Ethnicity and Age (3, 4) | | | | | | |
|--|--------|---------------------------|--|--|--|--|
| | Number | Anemia Low Hb/Hct % | | | | |
| Total All Race/Ethnic Groups | | | | | | |
| 6 - 11 Months | 98 | * | | | | |
| 12 - 17 Months | 124 | 11.3 | | | | |
| 18 - 23 Months | 49 | * | | | | |
| 24 - 35 Months | 94 | * | | | | |
| 36 - 59 Months | 185 | 8.6 | | | | |
| Total | 550 | 13.5 | | | | |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia Indicators by Race/Ethnicity and Age (3, 4) | | | | | | | | |
|--------------------------------|--|---------------------------|------------------|--------|---------------------------|--|--|--|--|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % | | | | |
| White, Not Hispanic | | | Asian | | | | | | |
| 6 - 11 Months | 15 | * | 6 - 11 Months | 0 | * | | | | |
| 12 - 17 Months | 32 | * | 12 - 17 Months | 0 | * | | | | |
| 18 - 23 Months | 13 | * | 18 - 23 Months | 0 | * | | | | |
| 24 - 35 Months | 18 | * | 24 - 35 Months | 0 | * | | | | |
| 36 - 59 Months | 77 | * | 36 - 59 Months | 1 | * | | | | |
| Total | 155 | 13.5 | Total | 1 | * | | | | |
| Black, Not Hispanic | | | Pacific Islander | | | | | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 1 1 | * | | | | |
| 12 - 17 Months | 4 | * | 12 - 17 Months | 3 | * | | | | |
| 18 - 23 Months | l i | * | 18 - 23 Months | 0 | * | | | | |
| 24 - 35 Months | 2 | * | 24 - 35 Months | 0 | * | | | | |
| 36 - 59 Months | 3 | * | 36 - 59 Months | 2 | * | | | | |
| Total | 10 | * | Total | 6 | * | | | | |
| Hispanic | | | Filipino | | | | | | |
| 6 - 11 Months | 9 | * | 6 - 11 Months | 0 | * | | | | |
| 12 - 17 Months | 18 | * | 12 - 17 Months | 0 | * | | | | |
| 18 - 23 Months | 3 | * | 18 - 23 Months | 0 | * | | | | |
| 24 - 35 Months | 13 | * | 24 - 35 Months | 1 | * | | | | |
| 36 - 59 Months | 29 | * | 36 - 59 Months | 0 | * | | | | |
| Total | 72 | * | Total | 1 | * | | | | |
| American Indian/Alaskan Native | | | All Other | | | | | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 1 | * | | | | |
| 12 - 17 Months | 3 | * | 12 - 17 Months | 2 | * | | | | |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 1 | * | | | | |
| 24 - 35 Months | 2 | * | 24 - 35 Months | 1 | * | | | | |
| 36 - 59 Months | 5 | * | 36 - 59 Months | 6 | * | | | | |
| Total | 11 | * | Total | 11 | * | | | | |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age

Children Aged < 5 Years (2) County: 018

Page:

Run Date:

6/3/09

| Anemia Indicators by Race/Ethnicity and Age (3, 4) | | | | | | | | |
|--|--------|---------------------------|--|--|--|--------|---------------------------|--|
| | Number | Anemia Low Hb/Hct % | | | | Number | Anemia Low Hb/Hct % | |
| Total All Race/Ethnic Groups | | | | | | | | |
| 6 - 11 Months | 27 | * | | | | | | |
| 12 - 17 Months | 62 | * | | | | | | |
| 18 - 23 Months | 18 | * | | | | | | |
| 24 - 35 Months | 37 | * | | | | | | |
| 36 - 59 Months | 123 | 10.6 | | | | | | |
| Total | 267 | 12.4 | | | | | | |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page:

Run Date:

6/3/09

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | Anemia | Indicators | y Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|---------------------------|---------------------------------|----------|---------------------------|
| | | Anemia Low Hb/Hct % | | <u> </u> | Anemia Low Hb/Hct % |
| | Number | 70 | | Number | 70 |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 24 | * | 6 - 11 Months | 31 | * |
| 12 - 17 Months | 19 | * | 12 - 17 Months | 53 | * |
| 18 - 23 Months | 12 | * | 18 - 23 Months | 19 | * |
| 24 - 35 Months | 14 | * | 24 - 35 Months | 34 | * |
| 36 - 59 Months | 48 | * | 36 - 59 Months | 98 | * |
| Total | 117 | 12.0 | Total | 235 | 21.7 |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 92 | * | 6 - 11 Months | 10 | * |
| 12 - 17 Months | 101 | 19.8 | 12 - 17 Months | 15 | * |
| 18 - 23 Months | 56 | * | 18 - 23 Months | 8 | * |
| 24 - 35 Months | 107 | 27.1 | 24 - 35 Months | 15 | * |
| 36 - 59 Months | 246 | 20.7 | 36 - 59 Months | 13 | * |
| Total | 602 | 22.6 | Total | 61 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 484 | 19.4 | 6 - 11 Months | 4 | * |
| 12 - 17 Months | 556 | 13.1 | 12 - 17 Months | 4 | * |
| 18 - 23 Months | 255 | 11.8 | 18 - 23 Months | 2 | * |
| 24 - 35 Months | 518 | 14.1 | 24 - 35 Months | 3 | * |
| 36 - 59 Months | 1,219 | 10.0 | 36 - 59 Months | 12 | * |
| Total | 3,032 | 12.9 | Total | 25 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 86 | * |
| 12 - 17 Months | 2 | * | 12 - 17 Months | 72 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 40 | * |
| 24 - 35 Months | 1 | * | 24 - 35 Months | 79 | * |
| 36 - 59 Months | 1 | * | 36 - 59 Months | 157 | 12.1 |
| Total | 4 | * | Total | 434 | 13.4 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Long Beach

Anemia Indicators by Race/Ethnicity and Age

Children Aged < 5 Years (2) County: 062

Page:

Run Date:

6/3/09

| | Anemia I | ndicators | y Race/E | thnicity an | d Age (3, | 4) |
|------------------------------|----------|---------------------------|----------|-------------|-----------|--------|
| | Number | Anemia Low Hb/Hct % | | | | Number |
| Total All Race/Ethnic Groups | | | | | | l |
| 6 - 11 Months | 731 | 19.6 | | | | |
| 12 - 17 Months | 822 | 14.0 | | | | |
| 18 - 23 Months | 392 | 14.5 | | | | |
| 24 - 35 Months | 771 | 16.3 | | | | |
| 36 - 59 Months | 1,794 | 12.2 | | | | |
| Total | 4,510 | 14.6 | | | | |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators | y Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|---------------------------|---------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 106 | 30.2 | 6 - 11 Months | 1 1 | * |
| 12 - 17 Months | 83 | * | 12 - 17 Months | 2 | * |
| 18 - 23 Months | 35 | * | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 63 | * | 24 - 35 Months | 2 | * |
| 36 - 59 Months | 122 | 13.1 | 36 - 59 Months | 2 | * |
| Total | 409 | 18.3 | Total | 8 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 11 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 6 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 7 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 7 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 23 | * | 36 - 59 Months | 1 | * |
| Total | 54 | * | Total | 1 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 1,153 | 25.8 | 6 - 11 Months | 1 | * |
| 12 - 17 Months | 1,300 | 19.9 | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 469 | 16.8 | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 1,251 | 17.2 | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 2,175 | 11.1 | 36 - 59 Months | 0 | * |
| Total | 6,348 | 17.2 | Total | 1 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 32 | * |
| 12 - 17 Months | 8 | * | 12 - 17 Months | 32 | * |
| 18 - 23 Months | 2 | * | 18 - 23 Months | 14 | * |
| 24 - 35 Months | 7 | * | 24 - 35 Months | 38 | * |
| 36 - 59 Months | 18 | * | 36 - 59 Months | 71 | * |
| Total | 36 | * | Total | 187 | 16.0 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia I | ndicators | 4) |
|------------------------------|----------|---------------------------|--------|
| | Number | Anemia Low Hb/Hct % | Number |
| Total All Race/Ethnic Groups | | | |
| 6 - 11 Months | 1,305 | 25.9 | |
| 12 - 17 Months | 1,431 | 19.6 | |
| 18 - 23 Months | 528 | 17.2 | |
| 24 - 35 Months | 1,368 | 17.2 | |
| 36 - 59 Months | 2,412 | 11.2 | |
| Гotal | 7.044 | 17.2 | |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | | | D (2.4) | | |
|--------------------------------|--------|----------------------|-------------------------------|--------|----------------------|
| | Anemia | Indicators | Race/Ethnicity and Age (3, 4) | | |
| | | Anemia Low Hb/Hct | | | Anemia Low Hb/Hct |
| | Number | % | | Number | % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 10 | * | 6 - 11 Months | 3 | * |
| 12 - 17 Months | 25 | * | 12 - 17 Months | 6 | * |
| 18 - 23 Months | 11 | * | 18 - 23 Months | 7 | * |
| 24 - 35 Months | 16 | * | 24 - 35 Months | 7 | * |
| 36 - 59 Months | 21 | * | 36 - 59 Months | 7 | * |
| Total | 83 | * | Total | 30 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 3 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 7 | * | 12 - 17 Months | 1 | * |
| 18 - 23 Months | 4 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 8 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 11 | * | 36 - 59 Months | 0 | * |
| Total | 33 | * | Total | 1 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 147 | 16.3 | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 284 | 11.3 | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 117 | 9.4 | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 200 | 11.0 | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 288 | 6.9 | 36 - 59 Months | 0 | * |
| Total | 1,036 | 10.5 | Total | 0 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 12 | * |
| 12 - 17 Months | 1 | * | 12 - 17 Months | 23 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 7 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 10 | * |
| 36 - 59 Months | 2 | * | 36 - 59 Months | 20 | * |
| Total | 3 | * | Total | 72 | * |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age

Children Aged < 5 Years (2) County: 021

Page:

Run Date:

6/3/09

| | Anemia I | ndicators |
|------------------------------|----------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 175 | 17.1 |
| 12 - 17 Months | 347 | 11.2 |
| 18 - 23 Months | 146 | 11.0 |
| 24 - 35 Months | 241 | 10.4 |
| 36 - 59 Months | 349 | 7.2 |
| Total | 1,258 | 10.7 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | A | | by Race/Ethnicity and Δαe (3, 4) | | |
|--------------------------------|--------|----------------------|----------------------------------|--------|----------------------|
| | Anemia | indicators | by Race/Ethnicity and Age (3, 4) | | |
| | | Anemia Low Hb/Hct | | | Anemia Low Hb/Hct |
| | Number | % | | Number | % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 17 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 6 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 7 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 6 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 13 | * | 36 - 59 Months | 0 | * |
| Total | 49 | * | Total | 0 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 0 | * |
| Total | 0 | * | Total | 0 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 2 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 2 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 3 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 3 | * | 36 - 59 Months | 0 | * |
| Total | 10 | * | Total | 0 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 1 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 1 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 2 | * | 36 - 59 Months | 4 | * |
| Total | 2 | * | Total | 6 | * |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Mariposa

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia I | ndicators |
|------------------------------|----------|---------------------------|
| | Number | Anemia Low Hb/Hct % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 20 | * |
| 12 - 17 Months | 9 | * |
| 18 - 23 Months | 10 | * |
| 24 - 35 Months | 6 | * |
| 36 - 59 Months | 22 | * |
| Total | 67 | * |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page:

Run Date:

6/3/09

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | Anemia | Indicators | y Race/Ethnicity and Age $^{(3,4)}$ | | |
|--------------------------------|--------|---------------------------|-------------------------------------|------------------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 18 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 31 | * | 12 - 17 Months | lő | * |
| 18 - 23 Months | 16 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 24 | * | 24 - 35 Months | Ö | * |
| 36 - 59 Months | 57 | * | 36 - 59 Months | 2 | * |
| Total | 146 | 8.9 | Total | 2 | * |
| Black, Not Hispanic | 1.10 | | Pacific Islander | - - | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | l o | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | o o | * |
| 24 - 35 Months | 1 | * | 24 - 35 Months | Ō | * |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 0 | * |
| Total | 1 | * | Total | 0 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 26 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 35 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 20 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 41 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 113 | 4.4 | 36 - 59 Months | 0 | * |
| Total | 235 | 10.6 | Total | 0 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 3 | * | 6 - 11 Months | 126 | 30.2 |
| 12 - 17 Months | 8 | * | 12 - 17 Months | 55 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 41 | * |
| 24 - 35 Months | 9 | * | 24 - 35 Months | 89 | * |
| 36 - 59 Months | 10 | * | 36 - 59 Months | 142 | 11.3 |
| Total | 30 | * | Total | 453 | 20.3 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Mendocino

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators |
|------------------------------|--------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 173 | 24.3 |
| 12 - 17 Months | 129 | 17.8 |
| 18 - 23 Months | 77 | * |
| 24 - 35 Months | 164 | 14.6 |
| 36 - 59 Months | 324 | 8.3 |
| Total | 867 | 15.3 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Run Date: 6/3/09

Page:

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|---------------------------|----------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 189 | 16.4 | 6 - 11 Months | 119 | 21.0 |
| 12 - 17 Months | 237 | 15.2 | 12 - 17 Months | 104 | 26.9 |
| 18 - 23 Months | 103 | 8.7 | 18 - 23 Months | 46 | * |
| 24 - 35 Months | 229 | 18.3 | 24 - 35 Months | 82 | * |
| 36 - 59 Months | 365 | 11.8 | 36 - 59 Months | 178 | 19.1 |
| Total | 1,123 | 14.3 | Total | 529 | 25.0 |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 41 | * | 6 - 11 Months | 3 | * |
| 12 - 17 Months | 61 | * | 12 - 17 Months | 4 | * |
| 18 - 23 Months | 38 | * | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 63 | * | 24 - 35 Months | 3 | * |
| 36 - 59 Months | 123 | 15.4 | 36 - 59 Months | 2 | * |
| Total | 326 | 26.4 | Total | 13 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 899 | 24.8 | 6 - 11 Months | 1 | * |
| 12 - 17 Months | 1,258 | 18.1 | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 592 | 17.7 | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 1,024 | 21.4 | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 1,900 | 14.2 | 36 - 59 Months | 1 | * |
| Total | 5,673 | 18.4 | Total | 2 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 210 | 26.2 |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 263 | 21.3 |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 49 | * |
| 24 - 35 Months | 2 | * | 24 - 35 Months | 150 | 24.7 |
| 36 - 59 Months | 2 | * | 36 - 59 Months | 282 | 19.1 |
| Total | 4 | * | Total | 954 | 22.4 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia I | ndicators |
|------------------------------|----------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 1,462 | 23.9 |
| 12 - 17 Months | 1,927 | 18.7 |
| 18 - 23 Months | 829 | 18.9 |
| 24 - 35 Months | 1,553 | 22.7 |
| 36 - 59 Months | 2,853 | 14.8 |
| Total | 8,624 | 19.0 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

County: 025

Page:

Run Date:

6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4 | 1) | |
|--------------------------------|--------|---------------------------|---------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 4 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 7 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 2 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 5 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 27 | * | 36 - 59 Months | 1 | * |
| Total | 45 | * | Total | 1 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 0 | * |
| Total | 0 | * | Total | 0 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 3 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 9 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 3 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 10 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 19 | * | 36 - 59 Months | 0 | * |
| Total | 44 | * | Total | 0 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 1 | * |
| 12 - 17 Months | 1 | * | 12 - 17 Months | 1 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 2 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 1 | * | 36 - 59 Months | 0 | * |
| Total | 2 | * | Total | 5 | * |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page:

Run Date:

6/3/09

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | Anemia I | ndicators |
|---|-----------------------|---------------------------|
| | Number | Anemia Low Hb/Hct % |
| Total All Race/Ethnic Groups 6 - 11 Months 12 - 17 Months 18 - 23 Months | 8 18 7 | * * |
| 24 - 35 Months 36 - 59 Months Total | 16 48 97 | * * |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page:

Run Date:

6/3/09

Children Aged < 5 Years (2)

Anemia Indicators by Race/Ethnicity and Age

| | | | | ` | |
|--------------------------------|-----------|---------------------------|---|-----------|---------------------------|
| | Anemia | Indicators | y Race/Ethnicity and Age ^{(3, 4} | ·) | |
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | 14dillooi | | Asian | Ttarribor | |
| 6 - 11 Months | 4 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 7 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 2 | * | 18 - 23 Months | 0 | * |
| | 5 | * | | 0 | * |
| 24 - 35 Months | 9 | * | 24 - 35 Months | | * |
| 36 - 59 Months | _ | * | 36 - 59 Months | | |
| Total | 27 | * | Total | 0 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 0 | * |
| Total | 0 | * | Total | 0 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 51 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 53 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 11 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 32 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 51 | * | 36 - 59 Months | 0 | * |
| Total | 198 | 19.2 | Total | 0 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 1 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 1 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 1 | * | 36 - 59 Months | 2 | * |
| Total | 1 | * | Total | 5 | * |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

County: 026

| | Anemia I | ndicators | | e/Ethnicity and Age (3, 4) | |
|-----------------------------|----------|----------------------|--------|----------------------------|--|
| | | Anemia Low Hb/Hct | | | |
| | Number | % | Number | | |
| otal All Race/Ethnic Groups | | | | | |
| 6 - 11 Months | 56 | * | | | |
| 12 - 17 Months | 61 | * | | | |
| 18 - 23 Months | 13 | * | | | |
| 24 - 35 Months | 38 | * | | | |
| 36 - 59 Months | 63 | * | | | |
| otal | 231 | 21.2 | | | |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|---------------------------|----------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | 1 | |
| 6 - 11 Months | 13 | * | 6 - 11 Months | 1 1 | * |
| 12 - 17 Months | 31 | * | 12 - 17 Months | 5 | * |
| 18 - 23 Months | 8 | * | 18 - 23 Months | 2 | * |
| 24 - 35 Months | 15 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 17 | * | 36 - 59 Months | 7 | * |
| Total | 84 | * | Total | 15 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 9 | * | 6 - 11 Months | 1 | * |
| 12 - 17 Months | 9 | * | 12 - 17 Months | 3 | * |
| 18 - 23 Months | 3 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 6 | * | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 6 | * | 36 - 59 Months | 1 | * |
| Total | 33 | * | Total | 6 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 1,041 | 22.5 | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 1,374 | 14.6 | 12 - 17 Months | 1 | * |
| 18 - 23 Months | 564 | 14.5 | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 794 | 10.2 | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 1,224 | 8.8 | 36 - 59 Months | 0 | * |
| Total | 4,997 | 14.1 | Total | 1 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 3 | * | 6 - 11 Months | 32 | * |
| 12 - 17 Months | 5 | * | 12 - 17 Months | 42 | * |
| 18 - 23 Months | 3 | * | 18 - 23 Months | 21 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 29 | * |
| 36 - 59 Months | 1 | * | 36 - 59 Months | 38 | * |
| Total | 12 | * | Total | 162 | 13.6 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Monterey

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators |
|------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 1,100 | 22.0 |
| 12 - 17 Months | 1,470 | 14.4 |
| 18 - 23 Months | 601 | 15.1 |
| 24 - 35 Months | 845 | 10.4 |
| 36 - 59 Months | 1,294 | 8.9 |
| Total | 5,310 | 14.1 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

County: 028

| | | | | | - |
|--------------------------------|--------|---------------------------|----------------------------------|--------|---------------------------|
| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 28 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 20 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 9 | * | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 11 | * | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 30 | * | 36 - 59 Months | 3 | * |
| Total | 98 | * | Total | 5 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 2 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 2 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 2 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 7 | * | 36 - 59 Months | 1 | * |
| Total | 14 | * | Total | 1 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 221 | 33.5 | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 132 | 32.6 | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 57 | * | 18 - 23 Months | 2 | * |
| 24 - 35 Months | 99 | * | 24 - 35 Months | 2 | * |
| 36 - 59 Months | 212 | 17.0 | 36 - 59 Months | 3 | * |
| Total | 721 | 28.8 | Total | 7 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 264 | 25.0 |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 44 | * |
| 18 - 23 Months | 2 | * | 18 - 23 Months | 71 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 72 | * |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 132 | 18.2 |
| Total | 3 | * | Total | 583 | 22.6 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age

Children Aged < 5 Years (2) County: 028

Page:

Run Date:

6/3/09

| | Anemia I | ndicators |
|------------------------------|----------|---------------------------|
| | Number | Anemia Low Hb/Hct % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 515 | 28.5 |
| 12 - 17 Months | 198 | 28.8 |
| 18 - 23 Months | 144 | 34.0 |
| 24 - 35 Months | 187 | 26.7 |
| 36 - 59 Months | 388 | 17.8 |
| Гotal | 1,432 | 26.0 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Δnemia | Indicators | by Race/Ethnicity and Age (3, 4) | ı | |
|--------------------------------|--------|---------------------------|----------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | I lace/Edimonty and Age | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 120 | 17.5 | 6 - 11 Months | l 1 | * |
| 12 - 17 Months | 56 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 13 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 66 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 154 | 11.0 | 36 - 59 Months | 1 | * |
| Total | 409 | 15.2 | Total | 2 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 1 | * | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 2 | * | 36 - 59 Months | 0 | * |
| Total | 3 | * | Total | 1 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 33 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 21 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 7 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 32 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 48 | * | 36 - 59 Months | 0 | * |
| Total | 141 | 9.2 | Total | 0 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 21 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 15 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 4 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 18 | * |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 16 | * |
| Total | 0 | * | Total | 74 | * |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

County: 029

Page:

Run Date:

6/3/09

| | Anemia | Indicators |
|----------------------------------|----------|---------------------------|
| | Number | Anemia Low Hb/Hct % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 175 | 15.4 |
| 12 - 17 Months 18 - 23 Months | 92 24 | * |
| 24 - 35 Months | 118 | 17.8 |
| 36 - 59 Months | 221 | 10.9 |
| Total | 630 | 13.5 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age

Children Aged < 5 Years (2) County: 030

Page:

Run Date:

6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4 |) | |
|--------------------------------|--------|---------------------------|---------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 182 | 13.7 | 6 - 11 Months | 247 | 19.8 |
| 12 - 17 Months | 206 | 7.3 | 12 - 17 Months | 203 | 9.9 |
| 18 - 23 Months | 75 | * | 18 - 23 Months | 54 | * |
| 24 - 35 Months | 186 | 9.7 | 24 - 35 Months | 225 | 14.2 |
| 36 - 59 Months | 319 | 9.4 | 36 - 59 Months | 516 | 10.9 |
| Total | 968 | 9.9 | Total | 1,245 | 12.8 |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 29 | * | 6 - 11 Months | 8 | * |
| 12 - 17 Months | 45 | * | 12 - 17 Months | 10 | * |
| 18 - 23 Months | 10 | * | 18 - 23 Months | 2 | * |
| 24 - 35 Months | 29 | * | 24 - 35 Months | 7 | * |
| 36 - 59 Months | 79 | * | 36 - 59 Months | 15 | * |
| Total | 192 | 15.1 | Total | 42 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 3,817 | 17.1 | 6 - 11 Months | 15 | * |
| 12 - 17 Months | 4,325 | 14.0 | 12 - 17 Months | 12 | * |
| 18 - 23 Months | 1,799 | 12.3 | 18 - 23 Months | 6 | * |
| 24 - 35 Months | 4,490 | 12.8 | 24 - 35 Months | 13 | * |
| 36 - 59 Months | 8,728 | 9.5 | 36 - 59 Months | 33 | * |
| Total | 23,159 | 12.4 | Total | 79 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 7 | * | 6 - 11 Months | 476 | 16.0 |
| 12 - 17 Months | 4 | * | 12 - 17 Months | 370 | 13.5 |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 112 | 16.1 |
| 24 - 35 Months | 8 | * | 24 - 35 Months | 423 | 15.6 |
| 36 - 59 Months | 6 | * | 36 - 59 Months | 696 | 12.4 |
| Total | 25 | * | Total | 2,077 | 14.3 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators |
|------------------------------|--------|---------------------------|
| | N | Anemia Low Hb/Hct % |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 4,781 | 17.0 |
| 12 - 17 Months | 5,175 | 13.6 |
| 18 - 23 Months | 2,058 | 12.2 |
| 24 - 35 Months | 5,381 | 12.9 |
| 36 - 59 Months | 10,392 | 9.7 |
| Total | 27,787 | 12.5 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|---------------------------|----------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 3 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 8 | * | 12 - 17 Months | 2 | * |
| 18 - 23 Months | 1 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 3 | * | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 9 | * | 36 - 59 Months | 10 | * |
| Total | 24 | * | Total | 13 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 17 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 31 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 12 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 32 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 48 | * | 36 - 59 Months | 0 | * |
| Total | 140 | 12.1 | Total | 0 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 69 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 101 | 11.9 | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 35 | * | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 100 | 19.0 | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 179 | 6.7 | 36 - 59 Months | 2 | * |
| Total | 484 | 14.0 | Total | 4 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 32 | * |
| 12 - 17 Months | 1 | * | 12 - 17 Months | 22 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 3 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 32 | * |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 32 | * |
| Total | 1 | * | Total | 121 | 11.6 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page:

Run Date:

6/3/09

Pasadena

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | Anemia I | ndicators |
|------------------------------|----------|---------------------------|
| | Number | Anemia Low Hb/Hct % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 121 | 26.4 |
| 12 - 17 Months | 165 | 13.9 |
| 18 - 23 Months | 52 | * |
| 24 - 35 Months | 169 | 14.8 |
| 36 - 59 Months | 280 | 6.1 |
| Total | 787 | 13.3 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|---------------------------|----------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 73 | * | 6 - 11 Months | 3 | * |
| 12 - 17 Months | 109 | 13.8 | 12 - 17 Months | 1 | * |
| 18 - 23 Months | 27 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 86 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 147 | 5.4 | 36 - 59 Months | 2 | * |
| Total | 442 | 11.3 | Total | 6 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 4 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 3 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 8 | * | 36 - 59 Months | 1 | * |
| Total | 16 | * | Total | 2 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 137 | 22.6 | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 173 | 17.3 | 12 - 17 Months | 1 | * |
| 18 - 23 Months | 45 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 135 | 16.3 | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 232 | 14.7 | 36 - 59 Months | 0 | * |
| Total | 722 | 17.5 | Total | 1 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 200 | 11.5 |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 155 | 12.3 |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 47 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 160 | 11.9 |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 168 | 6.0 |
| Total | 0 | * | Total | 730 | 10.4 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age

Children Aged < 5 Years (2) County: 031

Page:

Run Date:

6/3/09

| | Anemia I | ndicators |
|------------------------------|----------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 414 | 16.7 |
| 12 - 17 Months | 443 | 14.7 |
| 18 - 23 Months | 120 | 14.2 |
| 24 - 35 Months | 384 | 13.3 |
| 36 - 59 Months | 558 | 9.9 |
| Total | 1,919 | 13.4 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|---------------------------|----------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 11 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 20 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 1 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 11 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 28 | * | 36 - 59 Months | 0 | * |
| Total | 71 | * | Total | 0 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 1 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 5 | * | 36 - 59 Months | 1 | * |
| Total | 7 | * | Total | 1 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 11 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 11 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 2 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 6 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 13 | * | 36 - 59 Months | 0 | * |
| Total | 43 | * | Total | 0 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 6 | * |
| 12 - 17 Months | 1 | * | 12 - 17 Months | 1 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 2 | * |
| 36 - 59 Months | 1 | * | 36 - 59 Months | 9 | * |
| Total | 2 | * | Total | 18 | * |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age

Children Aged < 5 Years (2) County: 032

Page:

Run Date:

6/3/09

| | Anemia I | ndicators | (3, 4) |
|-----------------------------|----------|----------------------|--------|
| | | Anemia Low Hb/Hct | |
| | Number | % | Number |
| otal All Race/Ethnic Groups | | | • |
| 6 - 11 Months | 29 | * | |
| 12 - 17 Months | 34 | * | |
| 18 - 23 Months | 3 | * | |
| 24 - 35 Months | 19 | * | |
| 36 - 59 Months | 57 | * | |
| otal | 142 | 8.5 | |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators | y Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|---------------------------|---------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 150 | 14.0 | 6 - 11 Months | 9 | * |
| 12 - 17 Months | 184 | 7.1 | 12 - 17 Months | 11 | * |
| 18 - 23 Months | 104 | 10.6 | 18 - 23 Months | 3 | * |
| 24 - 35 Months | 185 | 12.4 | 24 - 35 Months | 11 | * |
| 36 - 59 Months | 377 | 8.8 | 36 - 59 Months | 24 | * |
| Total | 1,000 | 10.1 | Total | 58 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 69 | * | 6 - 11 Months | 2 | * |
| 12 - 17 Months | 82 | * | 12 - 17 Months | 2 | * |
| 18 - 23 Months | 36 | * | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 128 | 24.2 | 24 - 35 Months | 3 | * |
| 36 - 59 Months | 231 | 20.8 | 36 - 59 Months | 6 | * |
| Total | 546 | 23.4 | Total | 14 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 1,181 | 20.8 | 6 - 11 Months | 3 | * |
| 12 - 17 Months | 1,819 | 15.2 | 12 - 17 Months | 3 | * |
| 18 - 23 Months | 863 | 17.4 | 18 - 23 Months | 4 | * |
| 24 - 35 Months | 2,036 | 18.0 | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 4,456 | 11.1 | 36 - 59 Months | 7 | * |
| Total | 10,355 | 14.8 | Total | 18 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 2 | * | 6 - 11 Months | 129 | 17.8 |
| 12 - 17 Months | 2 | * | 12 - 17 Months | 172 | 10.5 |
| 18 - 23 Months | 1 | * | 18 - 23 Months | 83 | * |
| 24 - 35 Months | 1 | * | 24 - 35 Months | 170 | 18.8 |
| 36 - 59 Months | 3 | * | 36 - 59 Months | 376 | 10.1 |
| Total | 9 | * | Total | 930 | 12.9 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Riverside

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia I | ndicators |
|------------------------------|----------|---------------------------|
| | Number | Anemia Low Hb/Hct % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 1,545 | 20.4 |
| 12 - 17 Months | 2,275 | 14.3 |
| 18 - 23 Months | 1,095 | 16.3 |
| 24 - 35 Months | 2,535 | 17.9 |
| 36 - 59 Months | 5,480 | 11.2 |
| Total | 12,930 | 14.6 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age

Children Aged < 5 Years (2) County: 034

Page:

Run Date:

6/3/09

| | Anemia | Indicators | y Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|------------|---------------------------------|--------|------------|
| | | Anemia | | | Anemia |
| | | Low Hb/Hct | | | Low Hb/Hct |
| | Number | % | | Number | % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 352 | 23.3 | 6 - 11 Months | 250 | 15.2 |
| 12 - 17 Months | 314 | 22.0 | 12 - 17 Months | 248 | 14.9 |
| 18 - 23 Months | 211 | 14.2 | 18 - 23 Months | 107 | 22.4 |
| 24 - 35 Months | 385 | 17.7 | 24 - 35 Months | 262 | 22.5 |
| 36 - 59 Months | 699 | 14.7 | 36 - 59 Months | 586 | 16.0 |
| Total | 1,961 | 18.0 | Total | 1,453 | 17.3 |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 321 | 19.9 | 6 - 11 Months | 9 | * |
| 12 - 17 Months | 349 | 24.1 | 12 - 17 Months | 5 | * |
| 18 - 23 Months | 146 | 32.2 | 18 - 23 Months | 2 | * |
| 24 - 35 Months | 385 | 24.2 | 24 - 35 Months | 7 | * |
| 36 - 59 Months | 654 | 17.9 | 36 - 59 Months | 12 | * |
| Total | 1,855 | 21.8 | Total | 35 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 950 | 23.8 | 6 - 11 Months | 9 | * |
| 12 - 17 Months | 1,081 | 21.9 | 12 - 17 Months | 5 | * |
| 18 - 23 Months | 584 | 15.9 | 18 - 23 Months | 4 | * |
| 24 - 35 Months | 984 | 21.3 | 24 - 35 Months | 3 | * |
| 36 - 59 Months | 2,058 | 14.5 | 36 - 59 Months | 11 | * |
| Total | 5,657 | 18.8 | Total | 32 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 20 | * | 6 - 11 Months | 1,348 | 18.1 |
| 12 - 17 Months | 13 | * | 12 - 17 Months | 1,227 | 16.6 |
| 18 - 23 Months | 14 | * | 18 - 23 Months | 753 | 18.3 |
| 24 - 35 Months | 20 | * | 24 - 35 Months | 1,202 | 18.0 |
| 36 - 59 Months | 29 | * | 36 - 59 Months | 2,440 | 13.8 |
| Total | 96 | * | Total | 6,970 | 16.3 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Sacramento

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia I | ndicators |
|------------------------------|----------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 3,259 | 20.4 |
| 12 - 17 Months | 3,242 | 19.6 |
| 18 - 23 Months | 1,821 | 18.5 |
| 24 - 35 Months | 3,248 | 20.0 |
| 36 - 59 Months | 6,489 | 14.7 |
| 「otal | 18,059 | 17.9 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

County: 035

Page:

Run Date:

6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|-----------|---------------------------|----------------------------------|-----------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | 110111001 | | Asian | 110111001 | |
| 6 - 11 Months | 9 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 13 | * | 12 - 17 Months | l ő | * |
| 18 - 23 Months | 12 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 13 | * | 24 - 35 Months | lő | * |
| 36 - 59 Months | 17 | * | 36 - 59 Months | 2 | * |
| Total | 64 | * | Total | 2 | * |
| | 64 | | | | |
| Black, Not Hispanic | | * | Pacific Islander | | * |
| 6 - 11 Months | 2 | * | 6 - 11 Months | | * |
| 12 - 17 Months | | * | 12 - 17 Months | | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 1 4 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 1 | | 36 - 59 Months | | |
| Total | 5 | * | Total | 2 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 100 | 19.0 | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 177 | 14.1 | 12 - 17 Months | 1 | * |
| 18 - 23 Months | 84 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 125 | 22.4 | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 204 | 20.1 | 36 - 59 Months | 0 | * |
| Total | 690 | 18.1 | Total | 1 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 11 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 10 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 5 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 11 | * |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 31 | * |
| Total | 0 | * | Total | 68 | * |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 17B

2008 Pediatric Nutrition Surveillance (1) San Benito

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators |
|------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 123 | 18.7 |
| 12 - 17 Months | 203 | 12.8 |
| 18 - 23 Months | 101 | 13.9 |
| 24 - 35 Months | 150 | 21.3 |
| 36 - 59 Months | 255 | 18.0 |
| Total | 832 | 16.9 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page:

Run Date:

6/3/09

San Bernardino

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | Anemia | Indicators | by Race/Ethnicity and Age $^{(3,4)}$ | | |
|--------------------------------|--------|---------------------------|--------------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 142 | 11.3 | 6 - 11 Months | 16 | * |
| 12 - 17 Months | 156 | 12.2 | 12 - 17 Months | 17 | * |
| 18 - 23 Months | 76 | * | 18 - 23 Months | 6 | * |
| 24 - 35 Months | 151 | 19.9 | 24 - 35 Months | 19 | * |
| 36 - 59 Months | 374 | 14.4 | 36 - 59 Months | 42 | * |
| Total | 899 | 14.8 | Total | 100 | 10.0 |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 108 | 32.4 | 6 - 11 Months | 2 | * |
| 12 - 17 Months | 106 | 25.5 | 12 - 17 Months | 3 | * |
| 18 - 23 Months | 55 | * | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 142 | 21.8 | 24 - 35 Months | 3 | * |
| 36 - 59 Months | 321 | 21.8 | 36 - 59 Months | 6 | * |
| Total | 732 | 23.6 | Total | 15 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 1,128 | 24.1 | 6 - 11 Months | 2 | * |
| 12 - 17 Months | 1,867 | 16.5 | 12 - 17 Months | 6 | * |
| 18 - 23 Months | 1,067 | 15.9 | 18 - 23 Months | 4 | * |
| 24 - 35 Months | 2,288 | 17.1 | 24 - 35 Months | 9 | * |
| 36 - 59 Months | 4,711 | 11.2 | 36 - 59 Months | 9 | * |
| Total | 11,061 | 15.1 | Total | 30 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 213 | 14.1 |
| 12 - 17 Months | 1 | * | 12 - 17 Months | 333 | 12.3 |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 215 | 8.8 |
| 24 - 35 Months | 1 | * | 24 - 35 Months | 399 | 15.5 |
| 36 - 59 Months | 3 | * | 36 - 59 Months | 815 | 8.0 |
| Total | 6 | * | Total | 1,975 | 11.0 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 17B

2008 Pediatric Nutrition Surveillance (1) San Bernardino

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

County: 036

| | Anemia | Indicators |
|------------------------------|--------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 1,612 | 22.2 |
| 12 - 17 Months | 2,489 | 16.1 |
| 18 - 23 Months | 1,424 | 15.2 |
| 24 - 35 Months | 3,012 | 17.2 |
| 36 - 59 Months | 6,281 | 11.5 |
| Total | 14,818 | 14.9 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

County: 037

Page:

Run Date:

6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|-----------|---------------------------|----------------------------------|----------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | 110111001 | | Asian | - Tunion | |
| 6 - 11 Months | 732 | 14.1 | 6 - 11 Months | 71 | * |
| 12 - 17 Months | 934 | 9.0 | 12 - 17 Months | 82 | * |
| 18 - 23 Months | 408 | 7.8 | 18 - 23 Months | 36 | * |
| 24 - 35 Months | 868 | 16.4 | 24 - 35 Months | 72 | * |
| 36 - 59 Months | 1.734 | 10.2 | 36 - 59 Months | 203 | 14.8 |
| | , - | | | | _ |
| Total | 4,676 | 11.5 | Total | 464 | 16.6 |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 190 | 17.9 | 6 - 11 Months | 9 | * |
| 12 - 17 Months | 320 | 14.4 | 12 - 17 Months | 17 | * |
| 18 - 23 Months | 138 | 15.9 | 18 - 23 Months | 5 | * |
| 24 - 35 Months | 259 | 21.2 | 24 - 35 Months | 12 | * |
| 36 - 59 Months | 554 | 16.1 | 36 - 59 Months | 22 | * |
| Total | 1,461 | 16.8 | Total | 65 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 3,986 | 16.4 | 6 - 11 Months | 46 | * |
| 12 - 17 Months | 5,102 | 12.1 | 12 - 17 Months | 57 | * |
| 18 - 23 Months | 2,305 | 10.5 | 18 - 23 Months | 24 | * |
| 24 - 35 Months | 4,422 | 15.6 | 24 - 35 Months | 35 | * |
| 36 - 59 Months | 8,414 | 10.6 | 36 - 59 Months | 80 | * |
| Total | 24,229 | 12.8 | Total | 242 | 12.0 |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 10 | * | 6 - 11 Months | 1,696 | 13.9 |
| 12 - 17 Months | 9 | * | 12 - 17 Months | 1,613 | 12.8 |
| 18 - 23 Months | 4 | * | 18 - 23 Months | 592 | 11.8 |
| 24 - 35 Months | 23 | * | 24 - 35 Months | 1,425 | 20.0 |
| 36 - 59 Months | 17 | * | 36 - 59 Months | 2,448 | 15.9 |
| Total | 63 | * | Total | 7,774 | 15.3 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page:

Run Date:

6/3/09

| | Anemia | Indicators |
|------------------------------|--------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 6,740 | 15.5 |
| 12 - 17 Months | 8,134 | 12.0 |
| 18 - 23 Months | 3,512 | 10.6 |
| 24 - 35 Months | 7,116 | 16.9 |
| 36 - 59 Months | 13,472 | 11.8 |
| Total | 38,974 | 13.3 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 17B 2008 Pediatric Nutrition Surveillance (1) San Francisco

Anemia Indicators by Race/Ethnicity and Age

Children Aged < 5 Years (2)

Page:

Run Date:

6/3/09

| | A ! - | l | v Bace/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|------------|---------------------------------|--------|------------|
| | Anemia | Indicators | y Race/Ethnicity and Age (3, 4) | | |
| | | Anemia | | | Anemia |
| | | Low Hb/Hct | | | Low Hb/Hct |
| | Number | % | | Number | % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 3 | * | 6 - 11 Months | 109 | 11.0 |
| 12 - 17 Months | 8 | * | 12 - 17 Months | 236 | 9.3 |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 42 | * |
| 24 - 35 Months | 4 | * | 24 - 35 Months | 189 | 6.9 |
| 36 - 59 Months | 5 | * | 36 - 59 Months | 363 | 8.5 |
| Total | 20 | * | Total | 939 | 8.4 |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 4 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 22 | * | 12 - 17 Months | 2 | * |
| 18 - 23 Months | 12 | * | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 19 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 33 | * | 36 - 59 Months | 5 | * |
| Total | 90 | * | Total | 8 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 51 | * | 6 - 11 Months | 4 | * |
| 12 - 17 Months | 152 | 21.1 | 12 - 17 Months | 6 | * |
| 18 - 23 Months | 28 | * | 18 - 23 Months | 2 | * |
| 24 - 35 Months | 123 | 18.7 | 24 - 35 Months | 6 | * |
| 36 - 59 Months | 161 | 8.1 | 36 - 59 Months | 5 | * |
| Total | 515 | 17.1 | Total | 23 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 13 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 23 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 6 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 24 | * |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 62 | * |
| Total | 0 | * | Total | 128 | 20.3 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 17B

2008 Pediatric Nutrition Surveillance (1) San Francisco

Page:

Run Date:

6/3/09

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | Anemia | Indicators |
|------------------------------|--------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 184 | 15.2 |
| 12 - 17 Months | 449 | 15.8 |
| 18 - 23 Months | 91 | * |
| 24 - 35 Months | 365 | 12.1 |
| 36 - 59 Months | 634 | 11.2 |
| Total | 1,723 | 13.3 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 17B 2008 Pediatric Nutrition Surveillance (1) San Joaquin

Page:

Run Date:

6/3/09

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | Anemia | Indicators | y Race/Ethnicity and Age (3, 4) | | |
|----------------------------------|--------|---------------------------|----------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White Net Hieronia | Number | ,,, | Asian | Number | |
| White, Not Hispanic | 100 | 20.0 | Asian | 36 | * |
| 6 - 11 Months | 195 | 20.0 12.8 | 6 - 11 Months | 57 | * |
| 12 - 17 Months | 74 | 12.0 | 12 - 17 Months | 34 | * |
| 18 - 23 Months 24 - 35 Months | 152 | 13.8 | 18 - 23 Months 24 - 35 Months | 68 | * |
| 36 - 59 Months | 322 | 9.9 | 36 - 59 Months | 132 | 17.4 |
| | | | | | |
| Total | 843 | 12.8 | Total | 327 | 17.7 |
| Black, Not Hispanic | | * | Pacific Islander | _ | * |
| 6 - 11 Months | 28 | * | 6 - 11 Months | 7 | * |
| 12 - 17 Months | 69 | * | 12 - 17 Months | 8 | * |
| 18 - 23 Months | 36 | * | 18 - 23 Months | 5 | * |
| 24 - 35 Months | 84 | | 24 - 35 Months | 6 | |
| 36 - 59 Months | 183 | 10.9 | 36 - 59 Months | 21 | • |
| Total | 400 | 13.3 | Total | 47 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 865 | 26.2 | 6 - 11 Months | 3 | * |
| 12 - 17 Months | 1,268 | 19.1 | 12 - 17 Months | 8 | * |
| 18 - 23 Months | 494 | 20.4 | 18 - 23 Months | 3 | * |
| 24 - 35 Months | 1,149 | 18.5 | 24 - 35 Months | 4 | * |
| 36 - 59 Months | 2,168 | 10.1 | 36 - 59 Months | 15 | * |
| Total | 5,944 | 16.9 | Total | 33 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 552 | 15.2 |
| 12 - 17 Months | 3 | * | 12 - 17 Months | 801 | 10.2 |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 256 | 12.9 |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 492 | 14.2 |
| 36 - 59 Months | 2 | * | 36 - 59 Months | 860 | 7.8 |
| Total | 6 | * | Total | 2,961 | 11.3 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 17B

2008 Pediatric Nutrition Surveillance (1) San Joaquin

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia l | ndicators |
|----------------------------------|--------------|---------------------------|
| | Number | Anemia Low Hb/Hct % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 1,592 | 21.5 |
| 12 - 17 Months | 2,409 902 | 15.3 16.9 |
| 18 - 23 Months 24 - 35 Months | 1,955 | 17.3 |
| 36 - 59 Months | 3,703 | 9.8 |
| Total | 10,561 | 14.8 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 17B 2008 Pediatric Nutrition Surveillance (1) San Luis Obispo

Page:

Run Date:

6/3/09

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | Anemia | Indicators | y Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|---------------------------|---------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 83 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 115 | 19.1 | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 46 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 54 | * | 24 - 35 Months | 3 | * |
| 36 - 59 Months | 166 | 12.7 | 36 - 59 Months | 3 | * |
| Total | 464 | 18.8 | Total | 6 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 6 | * | 6 - 11 Months | 1 | * |
| 12 - 17 Months | 9 | * | 12 - 17 Months | 1 | * |
| 18 - 23 Months | 2 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 3 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 6 | * | 36 - 59 Months | 0 | * |
| Total | 26 | * | Total | 2 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 257 | 26.8 | 6 - 11 Months | 1 | * |
| 12 - 17 Months | 333 | 18.0 | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 115 | 19.1 | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 187 | 19.8 | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 464 | 10.3 | 36 - 59 Months | 4 | * |
| Total | 1,356 | 17.4 | Total | 6 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 110 | 31.8 |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 112 | 20.5 |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 18 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 76 | * |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 122 | 17.2 |
| Total | 0 | * | Total | 438 | 21.7 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 17B

2008 Pediatric Nutrition Surveillance (1) San Luis Obispo

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

County: 040

Page:

Run Date:

6/3/09

| | Anemia I | ndicators |
|------------------------------|----------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 458 | 27.3 |
| 12 - 17 Months | 570 | 18.8 |
| 18 - 23 Months | 181 | 19.3 |
| 24 - 35 Months | 324 | 19.8 |
| 36 - 59 Months | 765 | 11.9 |
| Total | 2,298 | 18.4 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page:

Run Date:

6/3/09

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|----------|---------------------------|----------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White Not Hieronia | INGINIDO | | Asian | Number | |
| White, Not Hispanic | 19 | * | Asian | 2 | * |
| 6 - 11 Months | 14 | * | 6 - 11 Months | 2 | * |
| 12 - 17 Months | 7 | * | 12 - 17 Months | 2 | * |
| 18 - 23 Months | 14 | * | 18 - 23 Months | 4 | * |
| 24 - 35 Months | 17 | * | 24 - 35 Months | 7 | * |
| 36 - 59 Months | | | 36 - 59 Months | | |
| Total | 71 | * | Total | 17 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 16 | * | 6 - 11 Months | 16 | * |
| 12 - 17 Months | 16 | * | 12 - 17 Months | 25 | * |
| 18 - 23 Months | 6 | * | 18 - 23 Months | 10 | * |
| 24 - 35 Months | 10 | * | 24 - 35 Months | 14 | * |
| 36 - 59 Months | 18 | * | 36 - 59 Months | 18 | * |
| Total | 66 | * | Total | 83 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 671 | 15.5 | 6 - 11 Months | 7 | * |
| 12 - 17 Months | 639 | 13.5 | 12 - 17 Months | 2 | * |
| 18 - 23 Months | 303 | 7.6 | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 515 | 8.9 | 24 - 35 Months | 4 | * |
| 36 - 59 Months | 857 | 6.7 | 36 - 59 Months | 12 | * |
| Total | 2,985 | 10.6 | Total | 25 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 2 | * | 6 - 11 Months | 379 | 14.5 |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 356 | 11.2 |
| 18 - 23 Months | 1 | * | 18 - 23 Months | 114 | 8.8 |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 276 | 14.5 |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 447 | 10.3 |
| Total | 3 | * | Total | 1,572 | 12.2 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

San Mateo

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators |
|----------------------------------|--------------|---------------------------|
| | Number | Anemia Low Hb/Hct % |
| Total All Race/Ethnic Groups | Number | ,,, |
| 6 - 11 Months | 1,112 | 15.2 |
| 12 - 17 Months 18 - 23 Months | 1,054 443 | 12.8 8.1 |
| 24 - 35 Months | 837 | 10.9 |
| 36 - 59 Months | 1,376 | 8.1 |
| Total | 4,822 | 11.2 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page:

Run Date:

6/3/09

Santa Barbara

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4 |) | |
|--------------------------------|--------|---------------------------|---------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 103 | 16.5 | 6 - 11 Months | 3 | * |
| 12 - 17 Months | 81 | * | 12 - 17 Months | 3 | * |
| 18 - 23 Months | 31 | * | 18 - 23 Months | l ĭ | * |
| 24 - 35 Months | 80 | * | 24 - 35 Months | 4 | * |
| 36 - 59 Months | 140 | 12.1 | 36 - 59 Months | 14 | * |
| Total | 435 | 13.8 | Total | 25 | * |
| Black, Not Hispanic | 100 | 1010 | Pacific Islander | | |
| 6 - 11 Months | 23 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 14 | * | 12 - 17 Months | 1 | * |
| 18 - 23 Months | 5 | * | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 20 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 22 | * | 36 - 59 Months | 1 | * |
| Total | 84 | * | Total | 3 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 1,978 | 20.2 | 6 - 11 Months | 4 | * |
| 12 - 17 Months | 1,616 | 17.3 | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 584 | 14.7 | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 1,614 | 15.6 | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 3,235 | 10.8 | 36 - 59 Months | 6 | * |
| Total | 9,027 | 15.1 | Total | 11 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 5 | * | 6 - 11 Months | 506 | 16.0 |
| 12 - 17 Months | 2 | * | 12 - 17 Months | 328 | 13.7 |
| 18 - 23 Months | 2 | * | 18 - 23 Months | 93 | * |
| 24 - 35 Months | 2 | * | 24 - 35 Months | 392 | 10.7 |
| 36 - 59 Months | 4 | * | 36 - 59 Months | 625 | 8.5 |
| Total | 15 | * | Total | 1,944 | 12.0 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 17B

2008 Pediatric Nutrition Surveillance (1) Santa Barbara

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia I | ndicators | |
|-----------------------------|----------|----------------------|--------|
| | | Anemia Low Hb/Hct | |
| | Number | % | Number |
| otal All Race/Ethnic Groups | | | • |
| 6 - 11 Months | 2,622 | 19.1 | |
| 12 - 17 Months | 2,045 | 16.6 | |
| 18 - 23 Months | 718 | 14.3 | |
| 24 - 35 Months | 2,112 | 14.4 | |
| 36 - 59 Months | 4,047 | 10.5 | |
| otal | 11.544 | 14.5 | |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age

Children Aged < 5 Years (2) County: 043

Page:

Run Date:

6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age ^{(3, 4} | 1) | |
|--------------------------------|--------|---------------------------|--|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 57 | * | 6 - 11 Months | 191 | 15.2 |
| 12 - 17 Months | 31 | * | 12 - 17 Months | 179 | 8.9 |
| 18 - 23 Months | 13 | * | 18 - 23 Months | 96 | * |
| 24 - 35 Months | 40 | * | 24 - 35 Months | 149 | 18.8 |
| 36 - 59 Months | 69 | * | 36 - 59 Months | 363 | 7.7 |
| Total | 210 | 12.9 | Total | 978 | 10.8 |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 23 | * | 6 - 11 Months | 3 | * |
| 12 - 17 Months | 28 | * | 12 - 17 Months | 4 | * |
| 18 - 23 Months | 9 | * | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 26 | * | 24 - 35 Months | 6 | * |
| 36 - 59 Months | 32 | * | 36 - 59 Months | 10 | * |
| Total | 118 | 18.6 | Total | 24 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 1,536 | 16.7 | 6 - 11 Months | 12 | * |
| 12 - 17 Months | 1,179 | 13.7 | 12 - 17 Months | 14 | * |
| 18 - 23 Months | 672 | 11.5 | 18 - 23 Months | 5 | * |
| 24 - 35 Months | 1,415 | 11.6 | 24 - 35 Months | 8 | * |
| 36 - 59 Months | 2,426 | 6.3 | 36 - 59 Months | 19 | * |
| Total | 7,228 | 11.2 | Total | 58 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 3 | * | 6 - 11 Months | 388 | 14.2 |
| 12 - 17 Months | 4 | * | 12 - 17 Months | 385 | 21.0 |
| 18 - 23 Months | 2 | * | 18 - 23 Months | 173 | 10.4 |
| 24 - 35 Months | 6 | * | 24 - 35 Months | 417 | 13.4 |
| 36 - 59 Months | 14 | * | 36 - 59 Months | 658 | 8.8 |
| Total | 29 | * | Total | 2,021 | 13.3 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 17B

2008 Pediatric Nutrition Surveillance (1) Santa Clara

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

County: 043

| | Anemia | Indicators |
|------------------------------|--------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 2,213 | 16.2 |
| 12 - 17 Months | 1,824 | 15.0 |
| 18 - 23 Months | 971 | 10.7 |
| 24 - 35 Months | 2,067 | 12.4 |
| 36 - 59 Months | 3,591 | 7.1 |
| Total | 10,666 | 11.7 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | | | (0.4) | | |
|--------------------------------|--------|----------------------|----------------------------------|--------|----------------------|
| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
| | | Anemia Low Hb/Hct | | | Anemia Low Hb/Hct |
| | Number | % | | Number | % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 36 | * | 6 - 11 Months | 1 | * |
| 12 - 17 Months | 45 | * | 12 - 17 Months | 1 | * |
| 18 - 23 Months | 13 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 22 | * | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 20 | * | 36 - 59 Months | 0 | * |
| Total | 136 | 17.6 | Total | 3 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 1 | * |
| 12 - 17 Months | 4 | * | 12 - 17 Months | 1 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 1 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 2 | * |
| Total | 6 | * | Total | 4 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 255 | 26.7 | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 247 | 20.2 | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 104 | 15.4 | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 193 | 18.7 | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 283 | 8.5 | 36 - 59 Months | 0 | * |
| Total | 1,082 | 17.9 | Total | 1 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 83 | * |
| 12 - 17 Months | 1 | * | 12 - 17 Months | 27 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 12 | * |
| 24 - 35 Months | 1 | * | 24 - 35 Months | 68 | * |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 66 | * |
| Total | 2 | * | Total | 256 | 9.8 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Santa Cruz

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia I | ndicators |
|------------------------------|----------|---------------------------|
| | Number | Anemia Low Hb/Hct % |
| Total All Race/Ethnic Groups | Number | 70 |
| 6 - 11 Months | 377 | 23.1 |
| 12 - 17 Months | 326 | 19.9 |
| 18 - 23 Months | 130 | 15.4 |
| 24 - 35 Months | 286 | 15.7 |
| 36 - 59 Months | 371 | 7.3 |
| Total | 1,490 | 16.4 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators | y Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|---------------------------|---------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| Maria Maria | Number | ,,, | | Number | ,,, |
| White, Not Hispanic | 000 | 45.0 | Asian | 40 | * |
| 6 - 11 Months | 336 | 15.2 | 6 - 11 Months | 18 | * |
| 12 - 17 Months | 415 | 8.7 | 12 - 17 Months | 15 | * |
| 18 - 23 Months | 193 | 9.8 | 18 - 23 Months | 9 | · • |
| 24 - 35 Months | 385 | 7.3 | 24 - 35 Months | 16 | · • |
| 36 - 59 Months | 692 | 9.8 | 36 - 59 Months | 31 | • |
| Total | 2,021 | 10.0 | Total | 89 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 8 | * | 6 - 11 Months | 1 | * |
| 12 - 17 Months | 20 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 13 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 18 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 34 | * | 36 - 59 Months | 2 | * |
| Total | 93 | * | Total | 3 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 62 | * | 6 - 11 Months | 1 | * |
| 12 - 17 Months | 63 | * | 12 - 17 Months | 4 | * |
| 18 - 23 Months | 24 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 51 | * | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 136 | 8.8 | 36 - 59 Months | 4 | * |
| Total | 336 | 12.2 | Total | 10 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 29 | * | 6 - 11 Months | 61 | * |
| 12 - 17 Months | 23 | * | 12 - 17 Months | 62 | * |
| 18 - 23 Months | 8 | * | 18 - 23 Months | 23 | * |
| 24 - 35 Months | 28 | * | 24 - 35 Months | 59 | * |
| 36 - 59 Months | 46 | * | 36 - 59 Months | 135 | 12.6 |
| Total | 134 | 23.9 | Total | 340 | 14.1 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4 | 1) | |
|-------------------------------|--------|---------------------------|---------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 3 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 1 | * | 12 - 17 Months | 1 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 6 | * | 36 - 59 Months | 0 | * |
| Total | 10 | * | Total | 1 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 1 | * | 36 - 59 Months | 0 | * |
| Total | 1 | * | Total | 0 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 3 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | l 1 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 3 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 2 | * | 36 - 59 Months | 0 | * |
| Total | 9 | * | Total | 0 | * |
| American Indian/Alaskan Nativ | е | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 1 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 0 | * |
| Total | 0 | * | Total | 1 | * |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Sierra

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia I | ndicators |
|------------------------------|----------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 7 | * |
| 12 - 17 Months | 3 | * |
| 18 - 23 Months | 0 | * |
| 24 - 35 Months | 3 | * |
| 36 - 59 Months | 9 | * |
| Total | 22 | * |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

County: 047

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|---------------------------|----------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| | Number | /0 | | Number | /0 |
| White, Not Hispanic | | | Asian | _ | |
| 6 - 11 Months | 88 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 80 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 27 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 78 | * | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 118 | 9.3 | 36 - 59 Months | 1 | * |
| Total | 391 | 18.7 | Total | 2 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 5 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 1 | * | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 7 | * | 36 - 59 Months | 0 | * |
| Total | 14 | * | Total | 1 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 11 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 26 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 7 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 22 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 35 | * | 36 - 59 Months | 1 | * |
| Total | 101 | 5.9 | Total | 1 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 3 | * | 6 - 11 Months | 48 | * |
| 12 - 17 Months | 10 | * | 12 - 17 Months | 54 | * |
| 18 - 23 Months | 2 | * | 18 - 23 Months | 30 | * |
| 24 - 35 Months | 7 | * | 24 - 35 Months | 46 | * |
| 36 - 59 Months | 11 | * | 36 - 59 Months | 88 | * |
| Total | 33 | * | Total | 266 | 15.4 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Siskiyou

Anemia Indicators by Race/Ethnicity and Age

Children Aged < 5 Years (2) County: 047

Page:

Run Date:

6/3/09

| | Anemia l | ndicators |
|------------------------------|----------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 151 | 26.5 |
| 12 - 17 Months | 175 | 17.1 |
| 18 - 23 Months | 66 | * |
| 24 - 35 Months | 156 | 15.4 |
| 36 - 59 Months | 261 | 10.0 |
| Total | 809 | 16.1 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

County: 048

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4 | 4) | |
|--------------------------------|--------|---------------------------|---------------------------------|--------|----------------------|
| | T | Anemia Low Hb/Hct % | | | Anemia Low Hb/Hct |
| | Number | % | | Number | % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 49 | * | 6 - 11 Months | 6 | * |
| 12 - 17 Months | 54 | * | 12 - 17 Months | 12 | * |
| 18 - 23 Months | 29 | * | 18 - 23 Months | 10 | * |
| 24 - 35 Months | 47 | * | 24 - 35 Months | 6 | * |
| 36 - 59 Months | 82 | * | 36 - 59 Months | 11 | * |
| Total | 261 | 8.0 | Total | 45 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 57 | * | 6 - 11 Months | 1 | * |
| 12 - 17 Months | 74 | * | 12 - 17 Months | 3 | * |
| 18 - 23 Months | 38 | * | 18 - 23 Months | 2 | * |
| 24 - 35 Months | 77 | * | 24 - 35 Months | 4 | * |
| 36 - 59 Months | 141 | 11.3 | 36 - 59 Months | 11 | * |
| Total | 387 | 16.3 | Total | 21 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 256 | 17.2 | 6 - 11 Months | 2 | * |
| 12 - 17 Months | 352 | 12.8 | 12 - 17 Months | 10 | * |
| 18 - 23 Months | 180 | 7.8 | 18 - 23 Months | 5 | * |
| 24 - 35 Months | 314 | 10.8 | 24 - 35 Months | 7 | * |
| 36 - 59 Months | 542 | 6.5 | 36 - 59 Months | 11 | * |
| Total | 1,644 | 10.5 | Total | 35 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 3 | * | 6 - 11 Months | 42 | * |
| 12 - 17 Months | 2 | * | 12 - 17 Months | 94 | * |
| 18 - 23 Months | 1 | * | 18 - 23 Months | 47 | * |
| 24 - 35 Months | 2 | * | 24 - 35 Months | 57 | * |
| 36 - 59 Months | 2 | * | 36 - 59 Months | 128 | 6.3 |
| Total | 10 | * | Total | 368 | 12.2 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age

Children Aged < 5 Years (2) County: 048

Page:

Run Date:

6/3/09

| | Anemia I | ndicators |
|------------------------------|----------|---------------------------|
| | Number | Anemia Low Hb/Hct % |
| Total All Race/Ethnic Groups | Number | |
| 6 - 11 Months | 416 | 16.3 |
| 12 - 17 Months | 601 | 13.1 |
| 18 - 23 Months | 312 | 9.3 |
| 24 - 35 Months | 514 | 13.4 |
| 36 - 59 Months | 928 | 7.3 |
| Total | 2,771 | 11.3 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|-----------|---------------------------|----------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | Ttarribor | | Asian | Hamboi | |
| 6 - 11 Months | 40 | * | 6 - 11 Months | 2 | * |
| 12 - 17 Months | 27 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 14 | * | 12 - 17 Months 18 - 23 Months | 1 | * |
| | 24 | * | | Ö | * |
| 24 - 35 Months | 46 | * | 24 - 35 Months | 4 | * |
| 36 - 59 Months | | | 36 - 59 Months | | |
| Total | 151 | 17.2 | Total | 7 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 1 | * |
| 12 - 17 Months | 1 | * | 12 - 17 Months | 1 | * |
| 18 - 23 Months | 5 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 4 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 8 | * | 36 - 59 Months | 0 | * |
| Total | 19 | * | Total | 2 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 303 | 27.4 | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 300 | 20.7 | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 166 | 16.3 | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 240 | 16.3 | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 461 | 14.5 | 36 - 59 Months | 0 | * |
| Total | 1,470 | 18.9 | Total | 1 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 2 | * | 6 - 11 Months | 63 | * |
| 12 - 17 Months | 7 | * | 12 - 17 Months | 31 | * |
| 18 - 23 Months | 5 | * | 18 - 23 Months | 22 | * |
| 24 - 35 Months | 12 | * | 24 - 35 Months | 24 | * |
| 36 - 59 Months | 15 | * | 36 - 59 Months | 49 | * |
| Total | 41 | * | Total | 189 | 10.6 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age

Children Aged < 5 Years (2) County: 049

Page:

Run Date:

6/3/09

| | Anemia | Indicators |
|------------------------------|--------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 412 | 25.7 |
| 12 - 17 Months | 367 | 19.9 |
| 18 - 23 Months | 214 | 16.4 |
| 24 - 35 Months | 304 | 17.8 |
| 36 - 59 Months | 583 | 13.2 |
| Total | 1,880 | 18.4 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Stanislaus

Page:

Run Date:

6/3/09

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|---------------------------|----------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | , , , | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 522 | 21.1 | 6 - 11 Months | 38 | * |
| 12 - 17 Months | 457 | 21.4 | 12 - 17 Months | 32 | * |
| 18 - 23 Months | 183 | 17.5 | 18 - 23 Months | 10 | * |
| 24 - 35 Months | 376 | 20.2 | 24 - 35 Months | 25 | * |
| 36 - 59 Months | 825 | 13.8 | 36 - 59 Months | 59 | * |
| Total | 2,363 | 18.2 | Total | 164 | 26.8 |
| Black, Not Hispanic | , | | Pacific Islander | | |
| 6 - 11 Months | 65 | * | 6 - 11 Months | 6 | * |
| 12 - 17 Months | 43 | * | 12 - 17 Months | 10 | * |
| 18 - 23 Months | 26 | * | 18 - 23 Months | 3 | * |
| 24 - 35 Months | 52 | * | 24 - 35 Months | 6 | * |
| 36 - 59 Months | 111 | 27.9 | 36 - 59 Months | 9 | * |
| Total | 297 | 26.6 | Total | 34 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 1,541 | 26.0 | 6 - 11 Months | 2 | * |
| 12 - 17 Months | 1,666 | 19.9 | 12 - 17 Months | 3 | * |
| 18 - 23 Months | 654 | 22.3 | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 1,453 | 20.2 | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 3,007 | 15.1 | 36 - 59 Months | 5 | * |
| Total | 8,321 | 19.5 | Total | 11 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 4 | * | 6 - 11 Months | 309 | 34.6 |
| 12 - 17 Months | 6 | * | 12 - 17 Months | 261 | 29.5 |
| 18 - 23 Months | 2 | * | 18 - 23 Months | 53 | * |
| 24 - 35 Months | 6 | * | 24 - 35 Months | 211 | 30.8 |
| 36 - 59 Months | 11 | * | 36 - 59 Months | 467 | 21.0 |
| Total | 29 | * | Total | 1,301 | 27.7 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Stanislaus

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

County: 050

| | Anemia I | ndicators |
|------------------------------|----------|---------------------------|
| | Number | Anemia Low Hb/Hct % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 2,487 | 26.3 |
| 12 - 17 Months | 2,478 | 21.4 |
| 18 - 23 Months | 931 | 21.8 |
| 24 - 35 Months | 2,130 | 21.3 |
| 36 - 59 Months | 4,494 | 16.0 |
| Total | 12,520 | 20.4 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators | y Race/Ethnicity and Age (3, 4) |) | |
|--------------------------------|--------|----------------------|---------------------------------|--------|----------------------|
| | | Anemia Low Hb/Hct | | | Anemia Low Hb/Hct |
| | Number | % | | Number | % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 98 | * | 6 - 11 Months | 11 | * |
| 12 - 17 Months | 121 | 22.3 | 12 - 17 Months | 14 | * |
| 18 - 23 Months | 53 | * | 18 - 23 Months | 4 | * |
| 24 - 35 Months | 103 | 13.6 | 24 - 35 Months | 16 | * |
| 36 - 59 Months | 185 | 9.7 | 36 - 59 Months | 30 | * |
| Total | 560 | 13.6 | Total | 75 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 5 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 7 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 11 | * | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 12 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 13 | * | 36 - 59 Months | 0 | * |
| Total | 48 | * | Total | 1 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 204 | 11.3 | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 218 | 12.8 | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 93 | * | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 192 | 13.0 | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 350 | 4.6 | 36 - 59 Months | 0 | * |
| Total | 1,057 | 9.5 | Total | 1 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 45 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 45 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 31 | * |
| 24 - 35 Months | 1 | * | 24 - 35 Months | 45 | * |
| 36 - 59 Months | 5 | * | 36 - 59 Months | 81 | * |
| Total | 6 | * | Total | 247 | 18.2 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page:

Run Date:

6/3/09

Sutter

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | Anemia | Indicators |
|------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 363 | 12.9 |
| 12 - 17 Months | 405 | 16.5 |
| 18 - 23 Months | 194 | 9.3 |
| 24 - 35 Months | 369 | 13.8 |
| 36 - 59 Months | 664 | 8.0 |
| Total | 1,995 | 11.8 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

County: 052

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|---------------------------|----------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 44 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 38 | * | 12 - 17 Months | l i | * |
| 18 - 23 Months | 15 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 52 | * | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 93 | * | 36 - 59 Months | 0 | * |
| Total | 242 | 21.5 | Total | 2 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 2 | * | 12 - 17 Months | Ō | * |
| 18 - 23 Months | 1 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 2 | * | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 1 | * | 36 - 59 Months | 0 | * |
| Total | 6 | * | Total | 1 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 40 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 36 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 26 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 61 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 169 | 10.1 | 36 - 59 Months | 0 | * |
| Total | 332 | 16.0 | Total | 0 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 208 | 16.3 |
| 12 - 17 Months | 1 | * | 12 - 17 Months | 103 | 8.7 |
| 18 - 23 Months | 1 | * | 18 - 23 Months | 68 | * |
| 24 - 35 Months | 3 | * | 24 - 35 Months | 159 | 10.7 |
| 36 - 59 Months | 6 | * | 36 - 59 Months | 281 | 2.8 |
| Total | 11 | * | Total | 819 | 8.8 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Tehama Anemia Indicators by Race/Ethnicity and Age Page: Run Date: 6/3/09

Children Aged < 5 Years (2) County: 052

| | Anemia I | ndicators |
|------------------------------|----------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 292 | 19.2 |
| 12 - 17 Months | 181 | 14.9 |
| 18 - 23 Months | 111 | 11.7 |
| 24 - 35 Months | 279 | 14.7 |
| 36 - 59 Months | 550 | 7.5 |
| Total | 1,413 | 12.6 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4 | -) | |
|--------------------------------|--------|---------------------------|---------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 6 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 4 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 8 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 15 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 23 | * | 36 - 59 Months | 2 | * |
| Total | 56 | * | Total | 2 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 1 1 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 1 | * | 36 - 59 Months | 0 | * |
| Total | 2 | * | Total | 0 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 3 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 5 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 1 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 4 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 13 | * | 36 - 59 Months | 0 | * |
| Total | 26 | * | Total | 0 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 3 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 10 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 5 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 8 | * |
| 36 - 59 Months | 2 | * | 36 - 59 Months | 15 | * |
| Total | 2 | * | Total | 41 | * |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia I | ndicators |
|------------------------------|----------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 12 | * |
| 12 - 17 Months | 19 | * |
| 18 - 23 Months | 14 | * |
| 24 - 35 Months | 28 | * |
| 36 - 59 Months | 56 | * |
| Total | 129 | 9.3 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age

Children Aged < 5 Years (2)

Page:

Run Date:

6/3/09

County: 054

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4 |) | |
|--------------------------------|--------|---------------------------|---------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | + | | Asian | | |
| 6 - 11 Months | 230 | 22.6 | 6 - 11 Months | 58 | * |
| 12 - 17 Months | 301 | 21.6 | 12 - 17 Months | 79 | * |
| 18 - 23 Months | 133 | 19.5 | 18 - 23 Months | 35 | * |
| 24 - 35 Months | 254 | 20.1 | 24 - 35 Months | 41 | * |
| 36 - 59 Months | 467 | 14.1 | 36 - 59 Months | 79 | * |
| Total | 1,385 | 18.8 | Total | 292 | 26.4 |
| Black, Not Hispanic | 1 | | Pacific Islander | | |
| 6 - 11 Months | 18 | * | 6 - 11 Months | 2 | * |
| 12 - 17 Months | 24 | * | 12 - 17 Months | 3 | * |
| 18 - 23 Months | 19 | * | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 21 | * | 24 - 35 Months | 2 | * |
| 36 - 59 Months | 77 | * | 36 - 59 Months | 5 | * |
| Total | 159 | 25.2 | Total | 13 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 2,177 | 24.8 | 6 - 11 Months | 5 | * |
| 12 - 17 Months | 3,017 | 19.8 | 12 - 17 Months | 9 | * |
| 18 - 23 Months | 1,358 | 17.6 | 18 - 23 Months | 4 | * |
| 24 - 35 Months | 2,455 | 20.1 | 24 - 35 Months | 7 | * |
| 36 - 59 Months | 5,193 | 13.9 | 36 - 59 Months | 20 | * |
| Total | 14,200 | 18.2 | Total | 45 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 8 | * | 6 - 11 Months | 259 | 23.6 |
| 12 - 17 Months | 16 | * | 12 - 17 Months | 377 | 18.8 |
| 18 - 23 Months | 8 | * | 18 - 23 Months | 167 | 19.8 |
| 24 - 35 Months | 18 | * | 24 - 35 Months | 293 | 19.8 |
| 36 - 59 Months | 45 | * | 36 - 59 Months | 575 | 14.8 |
| Total | 95 | * | Total | 1,671 | 18.4 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia | Indicators |
|------------------------------|--------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 2,757 | 24.6 |
| 12 - 17 Months | 3,826 | 19.8 |
| 18 - 23 Months | 1,725 | 18.1 |
| 24 - 35 Months | 3,091 | 20.5 |
| 36 - 59 Months | 6,461 | 14.3 |
| Total | 17,860 | 18.5 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

County: 055

| | A! - | | by Race/Ethnicity and Δαρ (3, | 4) | |
|--------------------------------|--------|--------------|-------------------------------|---------|--------------|
| | Anemia | Indicators | by Race/Ethnicity and Age (3, | <u></u> | |
| | | Anemia | | | Anemia |
| | | Low Hb/Hct % | | l., . | Low Hb/Hct % |
| | Number | % | | Number | % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 9 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 3 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 5 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 2 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 15 | * | 36 - 59 Months | 0 | * |
| Total | 34 | * | Total | 0 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 1 | * | 36 - 59 Months | 0 | * |
| Total | 1 | * | Total | 0 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 1 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 0 | * |
| Total | 1 | * | Total | 0 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 0 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 0 | * | 36 - 59 Months | 1 | * |
| Total | 0 | * | Total | 1 | * |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page:

Run Date:

6/3/09

Tuolumne

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | Anemia I | ndicators | nd Age ^(3, 4) | |
|------------------------------|----------|----------------------|--------------------------|-----------|
| | | Anemia Low Hb/Hct | | An Low |
| | Number | % | Number | _ |
| Total All Race/Ethnic Groups | | | • | |
| 6 - 11 Months | 9 | * | | |
| 12 - 17 Months | 3 | * | | |
| 18 - 23 Months | 6 | * | | |
| 24 - 35 Months | 2 | * | | |
| 36 - 59 Months | 17 | * | | |
| Total | 37 | * | | |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age

Children Aged < 5 Years (2) County: 056

Page:

Run Date:

6/3/09

| | Anemia | Indicators | y Race/Ethnicity and Age (3, 4 | ł) | |
|--------------------------------|--------|------------|--------------------------------|--------|------------|
| | | Anemia | | | Anemia |
| | | Low Hb/Hct | | | Low Hb/Hct |
| | Number | % | | Number | % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 168 | 17.3 | 6 - 11 Months | 9 | * |
| 12 - 17 Months | 219 | 11.4 | 12 - 17 Months | 13 | * |
| 18 - 23 Months | 74 | * | 18 - 23 Months | 3 | * |
| 24 - 35 Months | 175 | 10.3 | 24 - 35 Months | 3 | * |
| 36 - 59 Months | 277 | 7.6 | 36 - 59 Months | 15 | * |
| Total | 913 | 11.1 | Total | 43 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 13 | * | 6 - 11 Months | 3 | * |
| 12 - 17 Months | 28 | * | 12 - 17 Months | 1 | * |
| 18 - 23 Months | 11 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 21 | * | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 26 | * | 36 - 59 Months | 7 | * |
| Total | 99 | * | Total | 11 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 2,646 | 17.0 | 6 - 11 Months | 5 | * |
| 12 - 17 Months | 3,647 | 12.1 | 12 - 17 Months | 5 | * |
| 18 - 23 Months | 1,277 | 10.6 | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 2,780 | 11.0 | 24 - 35 Months | 4 | * |
| 36 - 59 Months | 4,716 | 7.3 | 36 - 59 Months | 5 | * |
| Total | 15,066 | 11.1 | Total | 20 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 2 | * | 6 - 11 Months | 167 | 15.6 |
| 12 - 17 Months | 0 | * | 12 - 17 Months | 263 | 6.5 |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 87 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 197 | 16.2 |
| 36 - 59 Months | 1 | * | 36 - 59 Months | 310 | 7.4 |
| Total | 3 | * | Total | 1,024 | 10.7 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Ventura

Page: Run Date: 6/3/09

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

| | Anemia I | ndicators |
|------------------------------|----------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 3,013 | 16.9 |
| 12 - 17 Months | 4,176 | 11.7 |
| 18 - 23 Months | 1,453 | 10.9 |
| 24 - 35 Months | 3,180 | 11.4 |
| 36 - 59 Months | 5,357 | 7.3 |
| Total | 17,179 | 11.1 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

County: 057

| | Anemia | Indicators | by Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|---------------------------|----------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 71 | * | 6 - 11 Months | 4 | * |
| 12 - 17 Months | 55 | * | 12 - 17 Months | 5 | * |
| 18 - 23 Months | 30 | * | 18 - 23 Months | 3 | * |
| 24 - 35 Months | 59 | * | 24 - 35 Months | 14 | * |
| 36 - 59 Months | 114 | 13.2 | 36 - 59 Months | 13 | * |
| Total | 329 | 16.4 | Total | 39 | * |
| Black, Not Hispanic | 1 020 | | Pacific Islander | 1 33 | |
| 6 - 11 Months | 2 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 4 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 6 | * | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 8 | * | 24 - 35 Months | 1 1 | * |
| 36 - 59 Months | 15 | * | 36 - 59 Months | 1 | * |
| Total | 35 | * | Total | 2 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 277 | 17.7 | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 248 | 14.1 | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 167 | 10.8 | 18 - 23 Months | 0 | * |
| 24 - 35 Months | 351 | 12.5 | 24 - 35 Months | 0 | * |
| 36 - 59 Months | 533 | 12.4 | 36 - 59 Months | 4 | * |
| Total | 1,576 | 13.5 | Total | 4 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 90 | * |
| 12 - 17 Months | 1 | * | 12 - 17 Months | 81 | * |
| 18 - 23 Months | 0 | * | 18 - 23 Months | 46 | * |
| 24 - 35 Months | 0 | * | 24 - 35 Months | 78 | * |
| 36 - 59 Months | 1 | * | 36 - 59 Months | 158 | 16.5 |
| Total | 3 | * | Total | 453 | 16.3 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page:

Run Date:

6/3/09

| | Anemia I | ndicators |
|------------------------------|----------|----------------------|
| | | Anemia Low Hb/Hct |
| | Number | % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months | 445 | 18.9 |
| 12 - 17 Months | 394 | 14.7 |
| 18 - 23 Months | 252 | 10.3 |
| 24 - 35 Months | 511 | 14.5 |
| 36 - 59 Months | 839 | 13.2 |
| Total | 2,441 | 14.5 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Run Date: 6/3/09

Page:

County: 058

| | Anemia | Indicators | Race/Ethnicity and Age (3, 4) | | |
|--------------------------------|--------|---------------------------|-------------------------------|--------|---------------------------|
| | Number | Anemia Low Hb/Hct % | | Number | Anemia Low Hb/Hct % |
| White, Not Hispanic | | | Asian | | |
| 6 - 11 Months | 139 | 15.8 | 6 - 11 Months | 10 | * |
| 12 - 17 Months | 125 | 20.0 | 12 - 17 Months | 17 | * |
| 18 - 23 Months | 68 | * | 18 - 23 Months | 9 | * |
| 24 - 35 Months | 121 | 18.2 | 24 - 35 Months | 13 | * |
| 36 - 59 Months | 257 | 13.2 | 36 - 59 Months | 40 | * |
| Total | 710 | 15.9 | Total | 89 | * |
| Black, Not Hispanic | | | Pacific Islander | | |
| 6 - 11 Months | 6 | * | 6 - 11 Months | 0 | * |
| 12 - 17 Months | 10 | * | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 4 | * | 18 - 23 Months | 3 | * |
| 24 - 35 Months | 12 | * | 24 - 35 Months | 2 | * |
| 36 - 59 Months | 13 | * | 36 - 59 Months | 0 | * |
| Total | 45 | * | Total | 5 | * |
| Hispanic | | | Filipino | | |
| 6 - 11 Months | 143 | 24.5 | 6 - 11 Months | 2 | * |
| 12 - 17 Months | 143 | 20.3 | 12 - 17 Months | 0 | * |
| 18 - 23 Months | 61 | * | 18 - 23 Months | 1 | * |
| 24 - 35 Months | 159 | 23.9 | 24 - 35 Months | 1 | * |
| 36 - 59 Months | 269 | 11.2 | 36 - 59 Months | 3 | * |
| Total | 775 | 18.5 | Total | 7 | * |
| American Indian/Alaskan Native | | | All Other | | |
| 6 - 11 Months | 1 | * | 6 - 11 Months | 28 | * |
| 12 - 17 Months | 3 | * | 12 - 17 Months | 19 | * |
| 18 - 23 Months | 4 | * | 18 - 23 Months | 10 | * |
| 24 - 35 Months | 1 | * | 24 - 35 Months | 22 | * |
| 36 - 59 Months | 3 | * | 36 - 59 Months | 54 | * |
| Total | 12 | * | Total | 133 | 14.3 |



⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Anemia Indicators by Race/Ethnicity and Age Children Aged < 5 Years (2)

Page: Run Date: 6/3/09

| | Anemia I | ndicators |
|----------------------------------|------------|---------------------------|
| | Number | Anemia Low Hb/Hct % |
| Total All Race/Ethnic Groups | | |
| 6 - 11 Months 12 - 17 Months | 329 317 | 19.8 20.2 |
| 18 - 23 Months 24 - 35 Months | 160 331 | 16.3 22.4 |
| 36 - 59 Months | 639 | 12.4 |
| Total | 1,776 | 17.3 |

⁽¹⁾ Reporting period is January 1 through December 31.(2) Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.